

# INDUCTION OF SYSTEMIC ACQUIRED DISEASE RESISTANCE IN PLANTS: POTENTIAL IMPLICATIONS FOR DISEASE MANAGEMENT IN URBAN FORESTRY

by Glynn C. Percival

**Abstract.** Systemic acquired resistance (SAR) is the phenomenon whereby a plant's own defense mechanisms are induced by prior treatment with either a biological or chemical agent. The concept of SAR has been widely recognized and studied for the past 100 years in relation to increasing resistance to fungal, bacterial, and viral pathogens of economically important crop plants. The use of SAR as part of a disease management strategy in relation to urban forestry has received little scientific investigation despite offering some potential for tree pathogen control. The objectives of this paper are to discuss the concepts of SAR and, where applicable, indicate how SAR may be incorporated into a disease management system by urban foresters. Areas discussed include historical background, persistence of SAR, mechanisms of SAR, induction of SAR by biological organisms and chemicals, interplanting with flowering woody plants to promote an SAR response, and SAR in woody plants.

**Key Words.** Induced resistance; integrated disease management; disease control; plant-pathogen interactions; fungicides; systemic acquired resistance; SAR; salicylic acid; isonicotinic acid; plant activator.

During their life cycle, urban trees are susceptible to a broad range of pathogenic fungi which, if uncontrolled, can result in high mortality rates. Infection can be via the roots (e.g., *Phytophthora cinnamomi* and *Armillaria mellea*, which interfere with the absorption of water and nutrients from the soil), via leaves (e.g., leaf spots [*Gnomonia platani*, *G. veneta*], which infect foliage and reduce photosynthesis), via flowers (e.g., fungal blights that attack the flowers and affect reproduction), and via fruit rots such as *Monilinia fructicola*, which interfere with reproduction and/

or storage of food reserves. Indeed, tree diseases can be so problematic as to limit the type of tree species that can grow in a large geographic area. The American chestnut (*Castanea dentata*) was annihilated in North America as a timber tree by chestnut blight (Dierauf et al. 1997), and the American elm is being eliminated as a shade tree by Dutch elm disease (Stipes and Campana 1981).

The problem of tree diseases can be further exacerbated because trees are often already weakened by the wide range of abiotic stresses they encounter within a town or city environment (e.g., root hypoxia, salinity, and drought). Trees in this weakened state are more susceptible to a range of pathogens that under normal circumstances do not pose a threat to their biology. For example, birch trees planted in the south of England tend to suffer from drought stress on an annual basis. This in turn makes them more susceptible to rust pathogens (*Cronartium* spp.). Such a problem is rarely encountered in Scotland where cooler, milder growing conditions prevail.

Presently, foresters tend to control these diseases primarily through the use of chemicals or by pruning diseased tissue and burning the arisings (British Standards Institute 1989). Tolerance to fungicides in, for example, the brown rot fungus or *Monilinia fructicola* (Agrios 1997); failure of fungicides to control diseases once a tree is infected, for example, by *Armillaria mellea*; increased legislative restrictions of the use of agrochemicals; and failure to achieve even distribution of chemicals throughout a tree canopy means that new techniques of pathogen control are required.

It is widely known that plants can defend themselves against pathogen infection through a variety of mechanisms that can be either local, constitutive, or inducible (Franceschi et al. 1998, 2000). Inducible resistance mechanisms such as systemic acquired resistance (SAR) are broad-spectrum plant defense responses that can be induced biologically by challenging a plant with a weaker strain of a specific pathogen or exposing a plant to natural and/or synthetic chemical compounds (Elliston et al. 1977). SAR has been studied by plant biologists for the past 100 years as a means to increase resistance to fungal, bacterial, and viral pathogens in crop plants such as potato, wheat, and rice (Agrios 1997).

The objectives of this review are to discuss the concepts of SAR to raise awareness of the potential and past uses of this technique to control plant diseases and, where applicable, highlight for urban foresters how SAR could be incorporated into a pathogen management system within towns and cities. It is important, however, to emphasize that SAR as a system to control tree diseases has received little scientific investigation (Karokene et al. 1999). Consequently, examples used throughout this paper rely heavily on results produced from herbaceous crop research.

### **HISTORICAL BACKGROUND**

The natural phenomenon of resistance development in response to pathogen infection was initially recognized by Ray (1901) and Beauvène (1901), working with *Botrytis cinerea* (gray mold). Beauvène (1899) had previously discovered that the virulence of a strain of *B. cinerea* could be varied by pre-exposing the pathogen to heat or cold. He then induced SAR in *Begonia*, by planting in soil previously inoculated with the heat- or cold-treated strains or by injecting inoculum directly into the plant. Regardless of the inoculation technique used, plant resistance developed to subsequent infections with highly virulent strains of the same fungus. The first controlled

laboratory study of SAR was performed by Ross (1961), who demonstrated that inoculation of a single leaf of tobacco with tobacco mosaic virus (TMV) reduced the severity of subsequent infections on other leaves. He coined the term "SAR" for the resistance that developed in the untreated portions of TMV-inoculated plants. Extending work on SAR to fungi, Cruickshank and Mandryk (1960) showed that high levels of resistance against *Peronospora tabacina* (blue mold) could be achieved within 3 weeks following inoculation in field-grown tobacco when a spore suspension of the same fungus was injected into stem tissue (Cohen and Kuc 1981).

Other recent, well-characterized examples of SAR have been recorded in both dicotyledonous and monocotyledonous plants (Hunt and Ryals 1996; Schneider et al. 1997; Mauch-Mani and Metraux 1998).

### **PERSISTENCE OF SAR**

Many diseases and decay processes in woody tissue develop over many years; however, the effect of inducing an SAR response upon retarding existing disease infection within a tree is unknown (Christiansen et al. 1999). Once induced, the SAR response in spruce (*Picea abies*) has been observed to persist for at least 1 year and possibly longer following inoculation, leading to the interesting hypothesis that a sublethal attack one year may strengthen rather than weaken a tree's defense (Christiansen, personal communication). Once induced, SAR can lead to long-lasting, broad-spectrum disease control. For example, inoculation of cucumber with *Pseudomonas lachrymans* or *Colletotrichum lagenarium* provided systemic resistance within a few days against 13 separate diseases, including those caused by fungi, bacteria, and viruses. A single SAR-inducing infection protected cucumber plants for 44 weeks, while a booster inoculation 2 to 3 weeks after the primary infection led to season-long, broad-spectrum resistance (Kuc and Richmond 1977). Whether such broad-spectrum resistance occurs in trees has

never been investigated. Further research in this area is now warranted.

### MECHANISMS OF SAR

Van Loon and Van Kammen (1970) and Gianinazzi et al. (1970) showed that viral infection of tobacco induced the accumulation of a distinct set of proteins, called pathogenesis-related proteins (PR proteins). Ward et al. (1991) demonstrated that at least nine gene families were induced in uninfected leaves of inoculated plants; these gene families are now known as SAR genes. Several of these SAR gene products have direct antimicrobial activity or are closely related to classes of antimicrobial proteins. These include  $\beta$ -1,3-glucanases, chitinases (Busam et al. 1997), cysteine-rich proteins related to thaumatin, beta-(1,3)-glucanase, and the PR-1 proteins (Anfoka and Buchenauer 1997). Further corroboration for the involvement of SAR genes in resistance comes from a range of transgenic plant experiments using seedling material (Alexander et al. 1993). The set of SAR genes that are induced differs among plant species. In cucumber, a class-III chitinase is the most highly induced SAR gene, while in tobacco and *Arabidopsis*, PR-1 and NPR1 are the predominant genes expressed (Cao et al. 1997, 1998). Such differences between species may reflect different evolutionary or breeding constraints that have selected for the most effective SAR response to the specific pathogen to which a species is most subject to attack.

Morphological and biochemical changes in SAR-protected plants that then become infected include a significantly faster lignification response, which corresponds with an increase in peroxidase activity (Ajilan and Potter 1992). Other changes include increased glucose and fructose concentrations in systemic tissue (Chandra and Bhatt 1998), an accumulation of fungi-toxic  $\beta$ -ionone derivatives (Wyatt and Kuc 1992), induction of lipoxygenase (Staub et al. 1992), antimicrobial fatty acid derivatives (Namai et al. 1993), phenylalanine ammonia-lyase, phytoalexins (Elliston et al. 1977), and

hydroxyproline-rich glycoprotein (Raggi 1998). Within conifers, inducible defense systems include secondary resin production, synthesis of new phenolics, traumatic resin duct formation, and initiation of a wound periderm (Franceschi et al. 2000).

A schematic diagram of the SAR response, based on available literature, is presented in Figure 1. It is important to emphasize, however, that the exact nature of the long-distant systemic signal (systemin) has not yet been fully elucidated, and aspects of systemic signaling in SAR remain unknown (see Ward et al. 1991; Kessmann et al. 1994; Chen et al. 1994 for further information).

### SAR INDUCED BY BIOLOGICAL AGENTS

Although a range of organisms is commercially available for use as biocontrol agents, nearly all are based on a direct antibiotic principle. For an organism to be classified as an SAR activator, certain criteria need to be fulfilled (see section titled SAR Induced by Chemical Means). Consequently, whether the organisms discussed in this section can be classified as SAR activators is unknown and requires further research to confirm resistance-inducing activity. In all of these cases, therefore, mechanisms other than SAR may be involved in increasing resistance.

Kessmann et al. (1994) reported that an extract of *Rhynoutria sachalinensis* gave good control of powdery mildews and other crop pathogens. However, the active principle(s) are not known. Extracts from *Bacillus subtilis* have been reported to induce resistance in barley, especially against powdery mildew (Steiner et al. 1988), while an infiltration of *Penicillium janczewskii* or its culture filtrate into melon and cotton increased peroxidase activity (a response associated with SAR) and resulted in increased protection against *Rhizoctonia solani* and elimination of the incidence of dampening-off symptoms (Madi and Katan 1998).

Another interesting class of agents is plant growth-promoting rhizobacteria, which are able




Figure 1. SAR response in plants.

to protect plants against foliar pathogens when used as a seed soak treatment (Maurhoffer et al. 1994). This strategy offers an exciting potential since disease control and increased tree health can be combined in a single seed treatment with naturally occurring microorganisms. A report examined the induction of SAR by a *Pseudomonas fluorescens* biocontrol strain (Maurhoffer et al. 1994). They demonstrated that growing plants in soil inoculated with this bacterium induced symptoms of SAR in leaf tissue. This response was absent in plants grown in uninoculated soil, or in soil inoculated with a different non-SAR-inducing strain. Similarly, Wei et al. (1991) reported on strains of *Pseudomonas* able to protect cucumber against a range of diseases. Mercer and Kirk (1984a,b) reported a series of laboratory tests designed to select microorganisms that would be effective in reducing infection by decay fungi and their subsequent performance under long-term field conditions. From preliminary tests of 22 bacterial isolates, three isolates of *Bacillus* spp. were selected with antibiotic properties. Whether *Pseudomonas fluorescens* was one of the 22 bacterial isolates used was not reported; however, the use of *Pseudomonas fluorescens* as a dual SAR inducer and biocontrol agent may warrant further investigation. Inducing SAR by chemical means and simultaneously inoculating with *Bacillus* UW85, a non-SAR-inducing microbial biocontrol strain, to investigate how expression of SAR affected the efficacy of this biocontrol agent was undertaken by Chen et al. (1996). Interestingly, SAR reduced the mortality of tobacco seedlings inoculated with zoospores of fungal diseases but had no effect on the growth of UW85. Combination of SAR and treatment with UW85 resulted in additive suppression of disease development.

*Verticillium dahliae*, to be marketed as DutchTrig® (Arcadis, Inc., the Netherlands), is under current trial for the biocontrol of Dutch elm disease. Although promising, the practical use of microorganisms to induce SAR at present is most likely restricted to selected trees grown on a small scale

under controlled or glasshouse conditions and has useful implications for the tree- and shrub-growing industry. Interestingly, no commercial biological product with an SAR mode of action has been introduced into the market, although DutchTrig® offers a possibility for the future.

### **SAR INDUCED BY CHEMICALS**

Chemicals that are able to induce SAR would offer a number of advantages over current conventional techniques for disease control in trees. Even if only a portion of the canopy is sprayed, the SAR response would be transduced throughout the canopy to provide long-term resistance. Similarly, unlike the classical biological control concept based on a predator-prey situation, SAR retains its efficacy irrespective of adverse environmental conditions such as low temperatures.

According to Kessmann et al. (1994), three criteria need to be fulfilled before a chemical agent can be classified as an “activator” of the SAR response: 1) the treated plants are resistant to the same number and type of diseases as those plants in which SAR has been biologically induced, 2) the chemical used has no direct antimicrobial activity or can be converted by the tree into antimicrobial metabolites, and 3) the same pre-infectious biochemical processes are induced as recorded in plant tissues after biological induction of SAR.

### **Salicylic Acid (SA)**

Although salicylic acid (aspirin) is recognized as an inducer of PR protein accumulation and SAR resistance when sprayed onto plants, and it fulfills all of the criteria of an activator, there are a number of problems with using SA as an SAR-inducing chemical. Reports of SA-mediated resistance are restricted to effects in the treated tissue, indicating that SA does not translocate efficiently throughout the plants when applied exogenously (Enyedi and Raskin 1993). Similarly, SA which is deposited upon a leaf surface, is rapidly broken down—resulting in only a short-

term response. Finally, only a narrow safety margin separates the rates at which the compound is effective and the rate at which it becomes strongly phytotoxic. Because of these limitations, SA has not been considered a practical solution to disease control.

## INA

A far more promising chemical is 2,6-dichloroisonicotinic acid (CGA 41396) and its methylester (CGA 41397), both referred to as INA. These compounds were discovered as able to induce systemic resistance in plants (Jensen et al. 1998), which provided good protection against fungal and bacterial pathogens of crops under glasshouse and field conditions (Table 1). Importantly, whether applied as a foliar spray or root drench, the spectrum of SAR activity is identical to the resistance observed after local pre-infection with SAR-inducing biological agents.

Plant responses to INA and INA analogs include induction of  $\beta$ -1,3-glucanases, chitinases, 6-phosphogluconate-dehydrogenase, various phenylpropanoid-derived metabolites, accelerated lipid metabolism, and peroxidase(s) synthesis (Seguchi et al. 1992; Staub et al. 1992), resulting in high-level resistance against plant diseases within 2 days of application. Histological studies showed that plants treated with INA respond to infection by a single-cell necrosis at the site of attempted penetration. The few hyphae that managed to penetrate become surrounded

by a cluster of necrotic cells and fungal growth ceases (Madamanchi and Kuc 1991).

## Other Chemicals

A number of reports have claimed resistance-inducing activity for a range of chemicals or plant extracts. Although these compounds have failed to fulfill the criteria established for SAR activators, they still could be considered as part of an integrated disease management system.

Plants pre-treated with fosetyl-Al, metalaxyl (Ward 1984), and triazoles (Hauthal 1993) accumulate phytoalexins faster than nontreated plants. The use of triazoles to promote SAR may be of interest for urban foresters. Carbendazim is classified as a benzimidazole and forms part of the gel formulation Bayleton BM recommended for tree work under British Standard guidelines (British Standards Institute 1989). A range of fungitoxicants (furmecyclox, fenpropimorph, carbendazim, and triadimenol) applied as pruning wound treatments on apple trees following inoculation with *Coriolum versicolor* were investigated by Gendle and Clifford (1983). No SAR-inducing properties of these chemicals, however, were investigated, and in all cases mode of application was via small surface-area pruning wounds. Although speculative, if triazoles also induce SAR-type responses, application as a foliar spray to a larger surface area in conjunction with application to pruning wounds may prove more effective in disease control. Further research is warranted to confirm this hypothesis.

Benzo-(1,2,3)-thiadiazole-7-carbothioic acid S-methyl ester (BTH) and its derivatives induce SAR in a number of plants (Gorlach et al. 1996), while probenazole (Oryzemate®) is a systemic compound used successfully to protect rice against *Pyricularia oryzae* (Kato et al. 1984). Because of their tolerance and efficacy in crops, both compounds have been

**Table 1. Activity of GCA 41396 in field trials (after Kessmann et al. 1994).**

Host/pathogen	INA rate	% infected leaf area		
		Check	CGA 41396	Standard
Tobacco/ <i>Peronospora</i>	20g/hl	29	12	17 <sup>a</sup>
Pear/ <i>Erwinia</i>	25g/hl	45	18	4 <sup>b</sup>
Pepper/ <i>Xanthomonas</i>	25g/hl	66	10	65 <sup>c</sup>
Rice/ <i>Pyricularia</i>	1kg/hl	12	3.5	0.5 <sup>d</sup>
Rice/ <i>Xanthomonas</i>	2kg/hl	8.2	0.2	0.1 <sup>e</sup>

Application was foliar, except in rice where the compound was applied into water.

Standards used: <sup>a</sup>Ridomil + mancozeb, 24 + 196g/hl; <sup>b</sup>streptomycin, 25g/hl; <sup>c</sup>copper + mancozeb, 200 + 100g/hl; <sup>d</sup>Oryzemate (probenazole) 2.4kg/ha; <sup>e</sup>TF-130, 2kg/ha.

commercialized, but, surprisingly their use in urban forestry remains untested.

The use of ethylene-producing substances, such as Ethrel (ethephon), as a foliar spray to cause a timed and coordinated shedding of leaves to reduce street sweeping and cleaning costs has been suggested by Moore (1998). Similarly, such a substance could be used to manipulate a timed ripening and shedding of fruits that are undesirable or pose a potential hazard within an urban context. The ethylene-releasing agent ethephon has been shown to induce SAR gene expression (Kessmann et al. 1994). Any compound that can be used to manipulate plant growth and simultaneously promote an SAR response would obviously be of great benefit within an urban forestry situation.

Applications of low concentrations of jasmonic acid (JA) to plants induce proteinase inhibitors, proline-rich cell wall protein, and a range of enzymes involved in plant defense reactions (Sticher et al. 1997). Additionally, significant *in vitro* activity of JA has been reported (Schweizer et al. 1993). Use of JA within an urban forestry program may, however, have limitations since JA is also recognized as a growth regulator affecting a range of physiological processes including root growth, senescence of leaves, and stomatal opening (Moore 1998).

Other chemical or plant extract agents with proposed SAR-inducing activity include various inorganic salts, compost and compost water extracts, low molecular weight proteins (elicitors) silicon, oxalate, phosphate, 2-thiouracil, polyacrylic acid, nucleic acids, unsaturated fatty acids and N-Trimethyl-L-lysine (Kessmann et al. 1994).

#### **INTERPLANTING WITH FLOWERING WOODY PLANTS**

As mentioned above, spraying plants with JA or methyl jasmonate can induce an SAR response but presents a number of undesirable side effects. A novel route to immunization against tree pathogens, however, may be to continually expose plants to JA vapor. Indeed, horticulturists

and amateur gardeners may already be using JA vapor to control plant pathogens without realizing it. Until recently, the concept of intercropping flowering plants within a glasshouse or small-scale field crop was believed to reduce pathogen and pest incidence as the flowering plant attracted a range of beneficial insects for pollination. It is now known that various species of *Artemisia*, for example, have high levels of JA in their leaves that slowly diffuse from leaf tissue into the atmosphere. Consequently, interspersing *Artemisia* within a crop of tomato resulted in induction of SAR and a significant reduction in disease incidence (Day 1993).

Planting an understory of woody plants (*Berberis*, *Pyracantha*, *Mahonia*, etc.) around a tree base is a recognized technique for suppressing weeds and simultaneously reducing the incidence of vandalism. The use of perennial flowering shrubs and/or short-lived flowering annuals as an understory may also provide a means of promoting an SAR response. Similarly, interplanting non-flowering trees with ornamental flowering species such as *Prunus* or *Crataegus* may induce an SAR response. With the exception of *Jasminum* spp., knowledge of flowering woody plants producing SAR-inducing chemicals within leaf, shoot, and flowering tissue is very limited but warrants further research. The results of a "desktop" literature search of plants possessing and on occasion diffusing JA from leaves, shoots, and flowers are shown in Table 2. Only plants suitable for use as an understory within existing tree plantings or those used by the landscape industry are presented.

Because JA is a naturally occurring growth regulator, this compound will undoubtedly be found in a further range of plants not listed in Table 2. Further research is required to determine other plants that release SAR-inducing chemicals, concentrations released by these plants, and influences of life cycle (flowering, seed set, etc.) upon rates of release, as well as concentrations required to induce the SAR re-

**Table 2. Plants with potential for inducing SAR.**

Plant	Author	Reference
<i>Jasminum</i> spp.	Day (1993)	New Scientist, January:36–40
<i>Artemisia</i> spp.	Day (1993)	New Scientist, January:36–40
<i>Tulip gesneriana</i> L.cvs Gudoshnik and Apeldoorn	Saniewski et al. (1998)	J. Plant Growth Reg. 17:179–183
<i>Populus</i> hybrids	Wilbert et al. (1998)	Analytical Biochem. 257:186–194
<i>Taxus canadensis</i> and <i>T. cuspidata</i>	Ketchum et al. (1999)	Biotech. Bioeng. 62:97–105
<i>Rosa</i> hybrids	Meir et al. (1998)	Postharvest Biol. Biotech. 13:235–243
Willow ( <i>Salix</i> spp.)	Constabel and Ryan (1998)	Phytochemistry 47: 507–511
Apple ( <i>Malus domestica</i> )	Fan et al. (1997)	J. Agric. Food Chem. 45:208–211
Sunflower	Bailly et al. (1992)	J. Plant Growth Reg. 11:349–355
Petunia	Tamari et al. (1995)	Physiol. Plantarium 94:45–50
Apple Golden Delicious	Perez et al. (1993)	J. Plant Growth Reg. 12:163–167
Olive	Sanz et al. (1993)	Phytochemistry 102(1):94–99

sponse in neighboring plants. Until further research provides answers to these questions, this technique, if feasible, is most likely restricted to trees grown on a small scale under controlled nursery or glasshouse conditions.

### SAR IN WOODY PLANTS

Very little literature exists on the use of SAR in controlling diseases of woody plants. Of that available, however, the bulk has been undertaken by Christiansen and colleagues at the Norwegian Forest Institute, Ås, Norway, using *Picea abies* as models against the fungal disease *Ceratocystis polonica* and the Eurasian spruce bark beetle (*Ips typographus*). In summary, their results have shown that at least two different mechanisms are involved in the induction of enhanced resistance of *P. abies*. These are that polyphenolic parenchyma cells in the bark respond actively to wounding, and particularly to fungal infection, with wounds and beetle attacks in the outer phloem effectively enveloped and rendered harmless by these cells. Traumatic resin ducts are formed in the xylem of trees surviving *I. typographus* attacks or mass inoculation with *C. polonica*. These resinous materials produced by epithelial cells lining these ducts may contain newly formed phenolic material fungistatic in

nature. Together the traumatic resin ducts and surrounding polyphenol-accumulating cells may provide a potent protective structure in which chemical substances accumulate and are ready to retard the longitudinal and inward spread of pathogenic organisms attempting invasion after wounding (Nagy et al. 2000). Induction of change in polyphenolic parenchyma cells were visible 6 days after induction, while traumatic duct formation required 2 weeks (Franceschi et al. 2000). Christiansen and Krokene (1999) concluded that “vaccination” with fungal inoculations can enhance the resistance of spruce trees to later beetle attack, although such “vaccination” of spruce trees is unlikely to have any practical application in forestry but may have potential for protecting valuable ornamental trees threatened by high bark beetle populations. Other work includes that of Okey and Sreenivan (1996), who recorded that application of SA as a seed treatment, spray, and soil drench, reduced the number of lesions, the size of the lesion area, and the percentage of necrotic leaf area of two clones of cacao infected by *Phytophthora palmivora*. Application of the SAR-inducing chemical INA as a foliar spray reduced fire blight (*Erwinia carotovora*) symptoms by 45% compared to noninoculated controls (Kessmann et al. 1994).

## CONCLUSIONS

Historically, emphasis in plant pathology has been placed on the discovery of new resistance genes by breeding and molecular techniques rather than on using the resistance potential already present in plants. Recent demonstrations of the effectiveness of SAR in laboratory and field situations presents interesting opportunities for the control of plant diseases by the urban forester. Experiments with crop plants have shown that SAR can lead to long-lasting, broad-spectrum disease control and can be used preventively to bolster general plant health. Ample evidence suggests that SAR is based on multiple natural defense mechanisms, and this makes it less likely that a pathogen can readily develop resistance to this control measure. The availability of this long-lasting, broad-spectrum and potentially stable solution to disease control may have an positive impact on urban forestry management.

## LITERATURE CITED

- Agrios, G.N. 1997. *Plant Pathology* (4th ed.). Academic Press, San Diego, CA.
- Ajilan, A.M., and D.A. Potter. 1992. Lack of effect of tobacco mosaic virus-induced systemic acquired resistance to anthropoid herbivores. *Phytopathology* 82:647-651.
- Alexander, D., R.M. Goodman, M. Gut-Rella, C. Glascock, and K. Weymann 1993. Increased tolerance to two Oomycete pathogens in transgenic tobacco expressing pathogenesis-related protein 1a. *Proc. Natl. Acad. Sci. USA* 90:7327-7331.
- Anfoka, G., and H. Buchenauer. 1997. Systemic acquired resistance in tomato against *Phytophthora infestans* by pre-inoculating with tobacco mosaic virus. *Physiol. Mol. Plant Pathol.* 50:85-101.
- Bailly, C., F. Corbineau, and D. Come. 1992. The effects of abscisic acid and methyl jasmonate on the 1-aminocyclopropane 1-carboxylic acid conversion to ethylene in hypocotyl segments of sunflower seedlings and their control by calcium and calmodulin. *Plant Growth Reg.* 11:349-355.
- Beauvène, J. 1899. Le *Botrytis cinerea* et la maladie de la toile. *Cr. Acad. Sci. Paris* 128:846-849.
- Beauvène, J. 1901. Essais d'immunization des vegetaux contre de maladies cryptogamiques. *Cr. Acad. Sci. Paris* 133:107-110.
- British Standards Institute. 1989. BS3998 Recommendations for Tree Work. British Standards Institute, London, England.
- Busam, G., H.H. Kassemeyer, and U. Matern, U. 1997. Differential expression of chitinases in *Vitis vinifera* L. responding to systemic acquired resistance activators or fungal challenge. *Plant Physiol.* 115:1029-1038.
- Cao, H., J. Glazebrook, J. Clarke, S. Volko, and X. Dong. 1997. The Arabidopsis NPR1 gene that controls systemic acquired resistance encodes a novel protein containing ankyrin repeats. *Cell* 88:57-63.
- Cao, H., X. Li, and X.N. Dong. 1998. Generation of broad spectrum disease resistance by over expression of an essential regulatory gene in systemic acquired resistance. *Proc. Nat. Acad. Sci. USA* 95:6531-6536.
- Chandra, A., and R.K. Bhatt. 1998. Biochemical and physiological response to salicylic acid in relation to the systemic acquired resistance. *Photosynthetica* 35:255-258.
- Chen, Z., H. Silva, and D. Klessig. 1994. Involvement of reactive oxygen species in the induction of systemic acquired resistance by salicylic acid in plants. *Science* 242:1883-1886.
- Chen, J., L.L. Jacobson, J. Handselman, and R.M. Goodman. 1996. Compatibility of systemic acquired resistance and microbial biocontrol for suppression of plant disease in a laboratory assay. *Mol. Ecol.* 5:73-80.
- Christiansen, E., P. Karokene, A.A. Berryman, V.R. Franceschi, T. Krekling, F. Lieutier, A. Lonneborg, and H. Solheim. 1999. Mechanical injury and fungal infection induce acquired resistance in Norway spruce. *Tree Physiol.* 19:399-403.
- Christiansen, E., and P. Karokene. 1999. Can Norway spruce trees be 'vaccinated' against attack by *Ips typographus*? *Agric. For. Entomol.* 1:185-187.
- Cohen, Y., and J. Kuc, J. 1981. Evaluation of systemic resistance to blue mold induced in tobacco leaves by prior stem inoculation with *Peronospora tabacina*. *Phytopathology* 71:783-787.
- Constabel, C.P., and C.A. Ryan. 1998. A survey of wound and methyl jasmonate-induced leaf polyphenol oxidase in crop plants. *Phytochemistry* 47: 507-511.
- Cruickshank, I.A.M., and M. Mandryk. 1960. The effect of stem infestations of tobacco with

- Peronospora tabacina* adam. on foliage reaction to blue mold. *J. Aust. Inst. Agric. Sci.* 26:369–372.
- Day, S. 1993. A shot in the arm for plants. *New Scientist* January:36–40.
- Dierauf, T., J. Artman, S. Elkins, L. Griffin, and G.J. Griffin. 1997. High level of chestnut blight control on grafted American chestnut trees inoculated with hypovirulent strains. *J. Arboric.* 23(2):87–88.
- Elliston, J., J. Kuc, E. Williams, and J. Raje. 1977. Relation of phytoalexin accumulation to local and systemic protection of bean against anthracnose. *Phytopathol. Z.* 88:114–130.
- Enyedi, A.J., and I. Raskin. 1993. Induction of UDP-glucose, salicylic acid glucosyl-transferase activity in tobacco mosaic virus-inoculated tobacco (*Nicotiana tabacum*) leaves. *Plant Physiol.* 101:1375–1380.
- Fan, X., J.P. Mattheis, J.K. Fellman, and M.E. Patterson. 1997. Effect of methyl jasmonate on ethylene and volatile production by summered apples depends on fruit developmental stage. *J. Agric. Food. Chem.* 45:208–211.
- Franceschi, V.R., T. Krekling, A.A. Berryman, and E. Christiansen. 1998. Specialized phloem parenchyma cells in Norway spruce (Pinaceae) are a primary site of defense reactions. *Am. J. Bot.* 85:601–615.
- Franceschi, V.R., P. Karokene, T. Krekling, and E. Christiansen. 2000. Phloem parenchyma cells are involved in local and distant defense responses to fungal inoculation or bark-beetle attack in Norway spruce (Pinaceae). *Am. J. Bot.* 87(3):314–326.
- Gendle, P., and D.R. Clifford. 1983. Movement, persistence and performance of fungitoxicants applied as pruning wound treatments on apple trees. *Ann. App. Biol.* 102:281–291.
- Gianinazzi, S., C. Martin, and J.C. Vallee. 1970. Hypersensibilite aux virus, temperature et proteines solubles chez le *Nicotiana Xanthi* n.c. Apparition de nouvelles macromolecules lors de la repression de la synthese virale. *Cr. Acad. Sci. D.* 270:2382–2386.
- Gorlach, J., S. Volrath, G. Knaufbeiter, G. Hengy, and U. Beckhove. 1996. Benzothiadiazole, a novel class of inducers of systemic acquired resistance, activates gene expression and disease resistance in wheat. *Plant Cell* 8: 629–643.
- Hauthal, H.G. 1993. *New Weg intelligenter Chemie: Beispiel Pflanzenschutz, Nachrichten aus Chemie. Tech. Lab.* 41:566–570.
- Hunt, M.D., and J.A. Ryals. 1996. Systemic acquired resistance signal transduction. *Crit. Rev. Plant Sci.* 15:583–606.
- Jensen, B.D., A.O. Latunedada, D. Hudson, and J.A. Lucas. 1998. Protection of *Brassica* seedlings against downy mildew and damping-off by seed treatment with CGA 245705, an activator of systemic acquired resistance. *Pestic. Sci.* 52:63–69.
- Karokene, P., E. Christiansen, H. Solheim, V.R. Franceschi, and A.A. Berryman. 1999. Induced resistance to pathogenic fungi in Norway spruce. *Plant Physiol.* 121:565–569.
- Kato, T., Y. Yamaguchi, T. Harano, T. Yokayama, and Y. Uyehara. 1984. Unsaturated hydroxy fatty acids, the self defensive substance in rice plants against rice blast disease. *Chem. Lett.* 12:409–412.
- Kessmann H., T. Staub, C. Hofmann, T. Maetzke, J. Herzog, E. Ward, S. Uknes, and J. Ryals. 1994. Induction of systemic acquired disease resistance in plants by chemicals. *Ann. Rev. Phytopathol.* 32:439–459.
- Ketchum, R.E.B., D.M. Gibson, R.B. Croteau, and M.L. Shuler. 1999. The kinetics of taxoid accumulation in cell suspension cultures of *Taxus* following elicitation with methyl jasmonate. *Biotech. Bioeng.* 62:97–105.
- Kuc, J., and S. Richmond. 1977. Aspects of the protection of cucumber against *Colletotrichum lagenarium* by *Colletotrichum lagenarium*. *Phytopathology* 67:533–536.
- Madamanchi, N.R., and J. Kuc. 1991. Induced systemic resistance in plants, pp 347–362. In Cole, G.T., and H.C. Hoch (Eds.). *The Fungal Spore and Disease Initiation in Plants and Animals*. Plenum, New York, NY.
- Madi, L., and J. Katan. 1998. *Penicillium janczewskii* and its metabolites, applied to leaves, elicit systemic acquired resistance to stem rot caused by *Rhizoctonia solani*. *Physiol. Mol. Plant Pathol.* 53:163–175.
- Mauch-Mani, B., and J.P. Mettraux. 1998. Salicylic acid and systemic acquired resistance to pathogen attack. *Ann. Bot.* 82:535–540.
- Maurhofer, M., C. Hase, J.P. Mettraux, and G. Defago. 1994. Induction of systemic resistance of tobacco to tobacco mosaic virus by the root colonizing *Pseudomonas fluorescens* strain CHA0: Influence of the *gacA* gene and pyoverdine production. *Phytopathology* 84:139–146.
- Meir, S., S. Droby, H. Davidson, S. Alsevia, L. Cohen, and B. Horev. 1998. Suppression of *Botrytis* rot in

- cut rose flowers by postharvest application of methyl jasmonate. *Postharvest Biol. Biotech.* 13:235–243.
- Mercer, P.C., and S.A. Kirk. 1984a. Biological treatments for the control of decay in tree wounds. I. Laboratory tests. *Ann. App. Biol.* 104:211–219.
- Mercer, P.C., and S.A. Kirk. 1984b. Biological treatments for the control of decay in tree wounds. II. Field tests. *Ann. App. Biol.* 104:221–229.
- Moore, G.M. 1998. Tree growth regulators: Issues of control, matters of management. *J. Arboric.* 24(1):10–19.
- Nagy, N.E., V.R. Franceschi, H. Solheim, T. Krekling, and E. Christiansen. 2000. Wound-induced traumatic resin duct development in stems of Norway spruce (*Pinaceae*): Anatomy and cytochemical traits. *Am. J. Bot.* 87(3):302–313.
- Namai, T., T. Kato, Y. Yamaguchi, and T. Hirukawa. 1993. Anti-rice blast activity and resistance induction of C-18 oxygenated fatty acids. *Biosci. Biotech. Biochem.* 57:611–613.
- Okey, E.N., and T.N. Sreenivan. 1996. Salicylic acid: A factor in systemic resistance of Caco to *Phytophthora palmivora*. *Proc. Brighton Crop. Prot. Conf.* 955–960.
- Perez, A.G., C. Sanz, D.G. Richardson, and J.M. Loias. 1993. Methyl jasmonate vapor promotes beta-carotene synthesis and chlorophyll degradation in golden delicious apple peel. *J. Plant Growth Reg.* 12:163–167.
- Raggi, V. 1998. Hydroxyproline-rich glycoprotein accumulation in TMV-infected tobacco showing systemic acquired resistance to powdery mildew. *J. Phytopathol.* 146:321–325.
- Ray, J. 1901. Les malaides cryptogamiques des vegetaux. *Rev. Gen. Bot.* 13:163–175.
- Ross, A.F. 1961. Localized acquired resistance to plant virus infection in hypersensitive hosts. *Virology* 14:329–339.
- Saniewski M., K. Miyamoto, and J. Ueda. 1998. Gum formation by methyl jasmonate in tulip shoots is stimulated by ethylene. *J. Plant Growth Reg.* 17:179–183.
- Sanz, L.C., J.C. Fernandez-Maculet, E. Gomez, B. Vioque, and J.M. Olias. 1993. Effect of methyl jasmonate on ethylene biosynthesis and stomatal closure in olive leaves. *Phytochemistry* 102(1):94–99.
- Schneider, M., P. Schweizer, P. Meuwly, and J.P. Mettraux. 1997. Systemic acquired resistance in plants. *Int. Rev. Cytol.* 168:303–340.
- Schweizer, P., R. Gees, and E. Mosinger. 1993. Effect of jasmonic acid on the interactions of barley (*Hordeum vulgare* L.) with the powdery mildew *Erysiphe graminis* f.sp. *hordei*. *Plant Physiol.* 102:503–511.
- Seguchi, K., M. Kurotaki, S. Sekido, and I. Yamaguchi. 1992. Action mechanism of N-cyanomethyl-2-chloroisonicotiana-mide in controlling rice blast disease. *J. Pestic. Sci.* 17:107–113.
- Staub, T., P. Ahl-Goy, and H. Kessmann. 1992. Chemically induced disease resistance in plants, pp 239–249. In Lyr, H., and Polter, C. *Proc. 10th Int. Symp. System. Fungic. Antifungal Compounds.*
- Steiner, U., E.C. Oerke, and F. Schonbeck. 1988. Zur Wirksamkeit der induzierten resistenz unter praktischen andbaubedingungen. *Z. Pflanzenkr. Pflanzenschutz* 95:506–517.
- Sticher, L., B. Mauch-Mani, and J.P. Mettraux, 1997. Systemic acquired resistance. *Ann. Rev. Phytopathol.* 35:235–270.
- Stipes, R.J., and R.J. Campana (Eds.). 1981. *Compendium of Elm Diseases.* American Phytopathological Society, St Paul, MN. 81 pp.
- Tamari, G., A. Borochoy, R. Atzorn, and U. Weiss. 1995. Methyl jasmonate induces pigmentation and flavanoid gene-transcription in petunia corollas—A possible role in wound response? *Physiol. Plantarum* 94:45–50.
- Van Loon, L.C., and A. Van Kammen. 1970. Polyacrylamide disc electrophoresis of the soluble proteins from *Nicotiana tabacum* var. Samsun and Samsun NN. II. Changes in protein constitution after infection with tobacco mosaic virus. *Virology* 40:199–211.
- Ward, E.W.B. 1984. Suppression of metalaxyl activity by glyphosphate: Evidence that host defense mechanisms contribute to metalaxyl inhibition of *Phytophthora megasperma* f. sp. *glycines* in soybeans. *Physiol. Plant Pathol.* 25:381–386.
- Ward, E.R., S.J. Uknes, S.C. Williams, S.S. Dincher, and D.L. Wiederhold. 1991. Coordinate gene activity in response to agents that induce systemic acquired resistance. *Plant Cell* 3:1085–1094.
- Wei, G., J. Kloepper, and S. Tuzun. 1991. Induction of systemic resistance in cucumber to *Colletotrichum orbiculare* by select strains of plant growth-promoting rhizobacteria. *Phytopathology* 81:1508–1512.

Wilbert, S.M., L.H. Ericsson, and M.P. Gordon. 1998. Quantification of jasmonic acid, methyl jasmonate and salicylic acid by capillary liquid chromatography electrospray tandem mass spectrometry. *Analytical Biochem.* 257:186–194.

Wyatt, S.E., and J. Kuc. 1992. The accumulation of  $\beta$ -ionone and 3-hydroxy esters of  $\beta$ -ionone in tobacco immunized by foliar inoculation with tobacco mosaic virus. *Phytopathology* 82:580–582.

*R.A. Barlett Tree Research Laboratory, Europe  
The University of Reading  
2 Early Gate  
Whiteknights  
Reading, U.K. RG6 6AU*

**Résumé.** La résistance systémique acquise est un phénomène par lequel les propres mécanismes de défense d'une plante sont déclenchés par des traitements préalables, soit avec un agent biologique ou chimique. Ce concept a été largement reconnu et étudié au cours des 100 dernières années, et ce en relation avec l'augmentation de la résistance aux champignons, aux bactéries et aux virus à caractère pathogénique chez des plantes de cultures importantes économiquement parlant. L'utilisation de ce concept en foresterie urbaine comme élément d'une stratégie de gestion des maladies a reçu peu d'intérêt de la part de la recherche scientifique, et ce en dépit que cette avenue puisse offrir un certain potentiel pour le contrôle des maladies des arbres. Les objectifs de cet article sont de discuter de ces concepts, là où ils sont applicables, ainsi que d'indiquer comment ce concept peut être incorporé par les forestiers urbains dans un système de gestion des maladies. Les aspects discutés incluent les connaissances historiques, la persistance de la résistance systémique, ses mécanismes, son déclenchement par des organismes biologiques et chimiques, son transfert dans les plantes ligneuses à fleurs, et sa présence chez les plantes ligneuses.

**Zusammenfassung.** Die systemisch erworbene Resistenz (SAR) ist das Phänomen, wobei der

pflanzeigene Abwehrmechanismus durch vorangehende Behandlung durch entweder ein biologisches oder chemisches Präparat hervorgerufen wird. Das Konzept von SAR ist weit bekannt und während der letzten 100 Jahre in Beziehung auf die wachsende Resistenz gegenüber fugalen, bakteriellen und virösen Pathogenen von wirtschaftlich genutzten Pflanzen untersucht worden. Der Gebrauch von SAR als Teil einer Managementstrategie in Bezug auf moderne Forstwirtschaft hat bislang wenig wissenschaftliche Forschung beansprucht, obwohl es ein Potential zur Kontrolle von Baumpathogenen bietet. Die Zielsetzung dieser Studie ist es, das Konzept von SAR zu diskutieren und, wo möglich, zu zeigen, wie SAR in ein Pflanzenschutzprogramm für Forstwirtschaft eingearbeitet werden kann. Die diskutierten Bereiche schließen historische Hintergründe, Persistenz von SAR, Mechanismen, Einführung von SAR durch Bioorganismen oder Chemikalien, Unterpflanzungen mit blühenden, holzigen Pflanzen, um eine SAR-Reaktion hervorzurufen, und SAR in holzigen Pflanzen.

**Resumen.** La resistencia sistémica adquirida (SAR, por sus siglas en inglés) es el fenómeno por el cual los mecanismos de defensa propios de las plantas son inducidos con tratamientos previos bien sea biológicos o químicos. El concepto de SAR ha sido ampliamente reconocido y estudiado en los pasados 100 años, con relación al incremento de la resistencia a patógenos fungosos, bacterianos o virales, económicamente importantes para las plantas. El uso de SAR como parte de una estrategia de manejo con relación a la dasonomía urbana, ha recibido poca atención científica, a pesar de ofrecer un potencial para el control de los patógenos en los árboles. Los propósitos de este reporte son discutir los conceptos de SAR y, donde es aplicable, indicar cómo puede ser incorporado el SAR a un sistema de manejo por los dasónomos urbanos. Los campos discutidos incluyen bagaje histórico, persistencia del SAR, mecanismos del SAR, inducción de SAR por organismos biológicos y químicos, mezcla con plantas leñosas con floración para promover una respuesta del SAR, y el estudio del mismo SAR en plantas leñosas.