

Effects of *Grosmannia clavigera* and *Leptographium longiclavatum* on Western White Pine Seedlings and the Fungicidal Activity of Alamo[®], Arbotect[®], and TREE-äge[®]

Stephen A. Wyka, Joseph J. Docola, Brian L. Strom, Sheri L. Smith, Douglas W. McPherson, Srđan G. Aćimović, and Kier D. Klepzig

Abstract. Bark beetles carry a number of associated organisms that are transferred to the host tree upon attack that are thought to play a role in tree decline. To assess the pathogenicity to western white pine (WWP; *Pinus monticola*) of fungi carried by the mountain pine beetle (MPB; *Dendroctonus ponderosae*), and to evaluate the potential for systemic prophylactic treatments for reducing fungal impacts, experiments were conducted with WWP seedlings to meet three objectives: 1) evaluate pathogenicity of two MPB-associated blue-stain fungi; 2) evaluate phytotoxicity of tree injection products; 3) evaluate the anti-fungal activity of tree injection products, *in vitro* and *in vivo*, toward the associated blue-staining fungi. To evaluate pathogenicity, seedlings were inoculated with *Grosmannia clavigera* or *Leptographium longiclavatum*, common fungal associates of MPB. Seedling mortality at four months after inoculation was 50% with *L. longiclavatum* and 90% with *G. clavigera*, both significantly higher than controls and thereby demonstrating pathogenicity. Phytotoxic effects of TREE-äge[®], Alamo[®], and Arbotect[®] were evaluated by stem injection; no phytotoxic effects were observed. Anti-fungal properties of the same three products were evaluated *in vitro* against *G. clavigera*, where Alamo was most active. Co-inoculation of *G. clavigera* and *L. longiclavatum* into seedlings after a stem injection of Alamo showed significantly less mortality and lesion formation than either species alone. Results support the hypothesis that MPB blue-stain associates, particularly *G. clavigera*, promote death of WWP when attacked by MPB. These findings suggest that the administration of a fungicide with insecticide for tree protection against bark beetles may be advantageous.

Key Words. Bark Beetles; Blue-Stain Fungi; Emamectin Benzoate; *Grosmannia clavigera*; *Leptographium longiclavatum*; Mountain Pine Beetle; *Pinus monticola*; Propiconazole; Systemic Fungicide; Systemic Insecticide; Thiabendazole; Tree Injection; Western White Pine.

Dendroctonus ponderosae, the mountain pine beetle (MPB), is an aggressive and destructive pest of many pine species in western North America. Lodgepole pine (*Pinus contorta* Douglas var. *latifolia* Engelman) is a favorite host, while MPB is also a serious pest of ponderosa pine, *P. ponderosa* Douglas ex C. Lawson, whitebark pine, *P. albicaulis* Engelman, western white pine, *P. monticola* Douglas ex D. Don, and limber pine, *P. flexilis* James (Tsuneda and Hiratsuka 1984; Yamaoka et al. 1990; Solheim and Krokene 1998; Carroll et al. 2003; Ono 2003; Lee et al. 2006a). Recently documented MPB outbreaks in western North America have resulted in millions of tree deaths (Carroll et al. 2003; Ono 2003) and reinvigorated interest in control options.

Adult MPBs mine the inner bark, moving from the attack site in an axial and planar orientation in

host trees. High attack densities and the successful establishment of brood, which mine laterally, result in tree girdling and subsequent host death. Successful reproduction by MPB depends on its relationship with associated fungi (Raffa and Berryman 1983; Yamaoka et al. 1990; Yamaoka et al. 1995; Six and Paine 1998; Rice et al. 2007b). The three main blue-stain ascomycetes isolated from *D. ponderosae* are *Grosmannia clavigera* (Robinson-Jeffery and Davidson) Zipfel, de Beer and Wingfield [= *Ophiostoma clavigerum* (Robinson-Jeffery and Davidson) Harrington], *Ophiostoma montium* (Rumbold) von Arx, and *Leptographium longiclavatum* Lee, Kim and Breuil (Tsuneda and Hiratsuka 1984; Yamaoka et al. 1995; Solheim and Krokene 1998; Kim et al. 2005; Lee et al. 2005; Lee et al. 2006b). These fungi are not only vectored by MPB but are also considered

to have a mutualistic relationship with this insect. It is suggested that these fungi aid the beetle in overwhelming host defenses (Raffa and Berryman 1983), providing nutrition (Six and Paine 1998; Adams and Six 2007), and perhaps a more favorable environment for beetle development (Lee et al. 2006a). For example, axenic MPBs are capable of entering and mining host trees, but offspring are apparently insufficiently nourished and attacks do not result in successful reproduction (Six and Paine 1998). In addition, fungal symbionts of MPBs have been shown to cause tree mortality by themselves (Mathre 1964; Basham 1970; Shrimpton 1973; Strobel and Sugawara 1986; Yamaoka et al. 1995). It appears that the combined action of the beetle and the fungi result in a more rapid host death (Amman et al. 1989; Yamaoka et al. 1995; Solheim and Krokene 1998; Lee et al. 2006a).

In a study using western white pine (WWP), *P. monticola*, trees *in situ* in northern California, U.S., TREE-äge® (4% [wt/wt] emamectin benzoate (EB), Syngenta Crop Protection, Greensboro, North Carolina, U.S.) injected trees had significantly reduced adult MPB gallery length compared to untreated trees (10.1 ± 2.1 and $120.6 \text{ cm} \pm 13.4 \text{ SE}$, respectively), and did not appear to be girdled by adult beetle activity (Strom et al. *unpublished*). However, significant tree mortality was observed in the study. To investigate potential causes of tree death, study trees were felled, wood samples collected, and fungal isolates cultured. The isolates were submitted for identification to Drs. Michael Wingfield and Wilhelm de Beer of the Forestry and Agricultural Biotechnology Institute (FABI), Pretoria, South Africa, who confirmed isolates were two species: *G. clavigera* and *L. longiclavatum*, both of which are common associates of MPB (Lee et al. 2005; Lee et al. 2006b).

In this study, the authors consider two factors that may impact tree survival: 1) injected formulation and 2) MPB fungal associates. Applied correctly, tree injections have been found to be both useful and effective against insects and pathogens. Doccola et al. (2011b) demonstrated that green ash (*Fraxinus pennsylvanica* Marsh.) successfully compartmentalized tree injection sites following treatment with TREE-äge for emerald ash borer (*Agrilus planipennis* Fairmaire). Furthermore, hemlock woolly adelgid (*Adelges tsugae* Annand)-infested eastern hemlock (*Tsuga canadensis* L.) recovered with new growth following imidacloprid tree injection (Doccola et al.

2012). Other researchers have not documented tree mortality from injections, and have generally found little or no phytotoxicity with currently available products when used according to label (Grosman et al. 2002; Smitley et al. 2010; Doccola et al. 2011a; Doccola et al. 2011b; McCullough et al. 2011).

Although Ophiostomatoid species vary in their pathogenicity to trees, a few do kill trees and this is especially true of species exotic to the host. Among these are *Ophiostoma novo-ulmi* (causal agent of Dutch elm disease), *Ceratocystis fagacearum* (Bretz) Hunt (causal agent of oak wilt), and *Raffaelea lauricola* (causal agent of laurel wilt). In addition, the native black stain root disease (*Leptographium wageneri*) can be a lethal vascular wilt pathogen of conifers in California, U.S. (Wagener and Mielke 1961). All of these species are vectored by insects that bore into trees. Furthermore, MPB-associated fungi were reported to cause injury and mortality following inoculations in lodgepole pine, jack pine, and lodgepole × jack hybrids (Strobel and Sugawara 1986; Yamaoka et al. 1990; Yamaoka et al. 1995; Lee et al. 2006b; Rice et al. 2007a; Rice et al. 2007b; Rice and Langor 2009). It is suggested that once fungal associates colonize the sapwood, they impede water and mineral transport in these tissues (Amman 1978). However, the authors are not aware of previous research regarding the virulence of *G. clavigera* and *L. longiclavatum* in *P. monticola*.

In this study, researchers report on the effects of systemic injections of TREE-äge, Alamo, and Arbotect to WWP seedlings and the subsequent challenge of these treatments via inoculations with *G. clavigera* or *L. longiclavatum*. The effects of the three systemic pesticides on the *in vitro* growth of *G. clavigera* were also investigated.

METHODS

Fungal Isolation

Trees cut from a field study in the Modoc National Forest, Alturas, California, U.S., were the source of wood samples used in the fungal isolations. Sample chips were removed from wooden block samples of TREE-äge-treated and untreated trees as described in Shigo (1986) using a No. 11/4 concave wood gouge. Chips were placed into Difco™ Malt Extract Agar (MEA) and incubated in the dark at 25°C for one week. Samples were then examined

for Ophiostomatoid fungi by observing characteristic conidiophores. From this isolation, two species were differentiated based on their morphology, *G. clavigera* by a more highly branched colony with more elongate and clavate shaped conidia, and *L. longiclavatum* by a less frequently branching colony with smaller and more cylindrical conidia (Six et al. 2003). Confirmatory identifications by DNA sequencing were conducted by Drs. Wingfield and de Beer. These isolates are now maintained in the culture collection (CMW) of FABI, under *G. clavigera* #38988 and *L. longiclavatum* #38989.

Fungicide Screening

Three systemic injection products including TREE-äge (4% [wt/wt] emamectin benzoate, Syngenta Crop Protection, Greensboro, North Carolina, U.S.), Alamo (14.3% [wt/wt] propiconazole, Syngenta Crop Protection, Greensboro, North Carolina, U.S.), and Arbotect (20% [wt/wt] thiabendazole, Syngenta Crop Protection, Greensboro, North Carolina, U.S.), were evaluated *in vitro* to determine their effect on growth of *G. clavigera*. To do this, serial dilutions of each treatment were established, from 10,000 to 1 ppm, along with a sterile water control. Individual filter papers, previously sterilized in an autoclave and left to dry in a sterile environment (1.5 cm, VWR, Radnor, Pennsylvania, U.S.), were dipped halfway into respective treatment solutions with sterile forceps and placed flat at the outer edge of an MEA agar plate, with the edge of the paper ~2 cm from the center of the plate (two filter papers per plate, three plates per treatment). Each plate medium was then streaked in the middle with conidia from a one-week old colony of *G. clavigera*. Researchers did not include *L. longiclavatum* in the screening because colonies did not survive cold storage. Plates were incubated at 25°C for one week, after which time measurements were made, and the experiment was concluded. The distance from the edge of each filter paper to the closest viable mycelium was measured (repression zone). If contamination obscured the view of the filter paper, no measurement was taken.

Mean repression of fungal mycelium was calculated in millimeters from six measured repression zones, two per Petri dish, for each treatment. Analysis of variance (ANOVA) for a hierarchical design with nested errors was implemented using

the MIXED procedure in SAS (V. 9.3, SAS Institute Inc., Cary, North Carolina, U.S.) with treatment as a fixed effect and dish within treatment as random, followed by comparisons among treatments using Fisher's protected LSD. A second analysis was performed on the subset of treatments obtained by eliminating the control to test for main and interaction effects of the factors fungicide and concentration. Demonstration of a highly significant fungicide by concentration interaction on repression was followed by orthogonal linear contrasts (via Estimate statements) to test for a response to increased concentration (on a log scale) of each fungicide.

SEEDLING EXPERIMENTAL OVERVIEW

Following the screening of fungicides *in vitro*, two experiments with seedlings were conducted. The objectives of the first experiment were to 1) test the pathogenicity of *G. clavigera* and *L. longiclavatum* in WWP seedlings, and 2) test the phytotoxicity of systemic pesticides in WWP seedlings. The objective of the second experiment was to test whether multiple inoculation points of fungi altered results compared to a single inoculation.

Experiment #1

Pathogenicity of fungal isolates

Pinus monticola seedlings were purchased from the University of Idaho, Franklin H. Pitkin Forest Nursery (Moscow, Idaho, U.S.), and potted into 15.24-cm pots using a 1:2:1 mix of compost, peat and perlite. From this group of seedlings (average basal diameter = 0.67 cm ± 0.01 SE and average height = 16.8 cm ± 0.5 SE), 50 individuals were selected and randomly assigned to 5 treatments (n=10 replicates per treatment). The five treatments were 1) *G. clavigera* inoculation, 2) *L. longiclavatum* inoculation, 3) co-inoculation of *G. clavigera* and *L. longiclavatum* after injection of systemic fungicide Alamo, 4) sterile MEA, or 5) de-ionized water. No seedlings were inoculated with a mix of both fungi only. It is known that these fungi are commonly associated and occur in common in nature. The single inoculations were designed to determine if a particular fungal species was more virulent than the other, while co-inoculations, which only followed seedling treatment with Alamo, were used to evaluate fungicidal effects

when fungi occurred in unison, as they would in the natural environment. Inoculations were carried out by first cleaning the outer bark with cotton rolls soaked in Clean-jet (25% isopropanol, 0.5% polyalkoxylated alcohol, Arborjet Inc., Woburn, Massachusetts, U.S.). A sterile razor blade was then used to make one slanting incision (~0.5 cm) into the stem. One full loop of inoculum was transferred from a one-week old fungal colony to each treated seedling using a flame-sterilized microbiological loop. After inoculation, the incision was closed and tightly sealed with Parafilm and aluminum foil to prevent contamination and desiccation and to immobilize the stem at the point of incision. For sterile agar and water controls, one loop full of sterile MEA or 100 μ L of sterile water, respectively, were placed into the stem incision and similarly sealed. The seedlings were kept in a greenhouse under 400 watt metal halide lamps (10-hour light cycle) and watered weekly. Assignment of seedlings to treatments and location of seedlings in the greenhouse was according to a completely randomized design. Study seedlings were assessed weekly for foliar necrosis and destructively autopsied by dissection after 120 days to document lesion development and to re-isolate fungal species from cross sections of seedling stems.

Phytotoxicity of fungicides

Another 40 seedlings were randomly assigned to four treatments to test the phytotoxicity of injected TREE-äge, Alamo, and Arbotect 20-S, relative to an untreated control (UTC). A Micro I.V. designed for this experiment (Arborjet, Inc., Woburn, Massachusetts, U.S.), based on a 10 cc capacity syringe fitted with 18 ga \times 2.5 cm syringe needle, administered TREE-äge, Alamo, or Arbotect by gravity feed into *P. monticola* seedlings. All treatments were infused by drip, where each seedling received 5 mL of a 10% solution. Injections were conducted by first cleaning the injection site using cotton rolls soaked in Clean-jet. A sterile razor blade was used to make a thin incision into the sapwood of the seedling, just large enough to allow a tight fit of the syringe tip. The syringe was then placed into the incision, while elastic horticultural tape was used to secure the site. The valve on the Micro I.V. was opened to allow the pesti-

cide to be administered by slow drip. The Micro I.V. was not removed until all of the solution had drained. The seedlings were arranged randomly in a separate section of the greenhouse from that which was occupied by the pathogenicity study, and so were held under the same environmental conditions previously described. Seedlings were visually inspected weekly for foliar phytotoxicity (e.g., foliar necrosis) and destructively autopsied after 120 days to document vascular health.

Autopsies and statistics

Seedlings were observed weekly to note symptoms of foliar necrosis and autopsies were conducted at 120 days post-treatment, earlier if all needles were dead, to assess cause of death (Table 1). Where oxidized or necrotic phloem completely encircled the stem, the seedling was categorized as dead. The earliest mortality was observed at 40 days in seedlings inoculated with *G. clavigera*. After 120 days, the percent circumferential lesion was measured in all remaining treatments. Circumferential lesion formation reflects the extent of stem girdling, a critical metric to tree survival. Autopsies were conducted using the following steps: 1) the bark was removed at the base of each seedling, up to the first node (first year stem growth); lesion circumference and height were measured; and 2) thin transverse sections of treatments were taken for microscopic examination of discoloration (oxidation) and occlusions of the vascular tissues.

Statistical analyses were carried out separately for each experiment. For each experiment, the null hypothesis that mortality was the same for all treatments was tested using Fisher's exact test (FET) on an Rx2 frequency table with seedlings classified as living or dead for each of R treatments (R = 5 or 4 for the pathogenicity and phytotoxicity studies, respectively). Rejection of the null hypothesis at significance level 0.05 was followed by pairwise comparisons of treatments using FET with Bonferroni-adjusted *P*-values on the corresponding 2x2 tables. For the pathogenicity study, the two control treatments were compared first and, when a non-significant outcome ($P > 0.10$) was observed; they were pooled for comparisons with other treatments (Table 1). One-way analysis of variance (ANOVA) was used

Table 1. Evaluation of mortality 120 days after inoculations and injection of *Pinus monticola* seedlings with respective treatments in Experiment 1. Mortality rates for each treatment are based on 10 seedlings.

Study	Treatment	% Mortality \pm SE	Foliar symptoms	Vasc. disc. ^z	Notes
Pathogenicity	<i>G. clavigera</i>	90 \pm 10.0% a ^y	desiccated	yes	vascular lesions
	<i>L. Longiclavatum</i>	50 \pm 16.7% ab	desiccated	yes	vascular lesions
	Alamo/B.S. ^x	10 \pm 10.0% b	desiccated	yes	vascular lesions
	Pooled controls	15 \pm 8.0% b			
	[100 μ L DI H ₂ O	20 \pm 13.0%	desiccated	no	dehydrated
	Sterile agar	10 \pm 10.0%	desiccated	no	dehydrated]
Phytotoxicity	TREE-äge	0.0 \pm 0.0% a	normal	no	n/a
	Alamo	0.0 \pm 0.0% a	normal	no	n/a
	Arbotect	0.0 \pm 0.0% a	normal	no	n/a
	UTC ^w	10 \pm 0.0% a	desiccated	yes	unidentified vascular infection

^z Vascular discoloration evident.

^y Treatments within a study with the same letter are not significantly different based on Fisher's exact test with Bonferroni-adjusted *P*-values.

^x Blue-stain (B.S.) both *G. clavigera* and *L. longiclavatum*.

^w Untreated Control (UTC).

to compare the differences in foliar browning, circumferential girdling, and axial lesion length. Lesion length was log-transformed prior to analysis to decrease variance heterogeneity. Pairwise comparisons among treatments were carried out with Tukey's HSD procedure at the 0.05 level of significance using the MIXED procedure in SAS.

Experiment #2

Dual inoculation of *G. clavigera*

As before, *P. monticola* seedlings (mean caliper and height of 0.5 \pm 0.02 and 15.0 cm \pm 0.4 SE respectively), were purchased from the University of Idaho, Franklin H. Pitkin Forest Nursery (Moscow, Idaho, U.S.). In this experiment, *G. clavigera* and sterile MEA agar, with 10 replicates per treatment were employed. *Leptographium longiclavatum* was not used in this experiment because it failed to survive cold storage between experiments. Inoculations were made as in Experiment 1, but with a second inoculation being made on the opposite side of the stem (180 degrees to the first) and offset by 1 cm in height. The lesion data from Experiment 1 were used to determine the offset distance between incisions in this experiment. Seedlings were kept in a greenhouse under the same environmental conditions as previously stated and destructively autopsied after 40 days. This was sufficient time to observe needle browning. Lesion development was not documented, as researchers were only concerned with tree mortality. Cross sections of stems with fungal inoculations were plated on MEA to recover fungal species.

RESULTS

Fungicide Screening

The effect of concentration on repression of *G. clavigera* growth *in vitro* (Table 2) was not the same for all fungicides ($F_{8,29} = 17.76$, $P < 0.001$). There was no evidence of repression with increasing concentrations of TREE-äge (from 1 to 10,000 ppm, $t_{31} = 0.05$, $P = 0.962$). In contrast, there was significant repression observed with increasing concentrations of the fungicide treatments Alamo and Arbotect ($t_{31} = 15.46$, $P < 0.001$ and $t_{31} = 10.28$, $P < 0.001$, respectively). Treatments of Alamo at 1,000 and 100 ppm, showed significantly greater repression than respective concentrations of Arbotect ($P < 0.05$) (Table 2). However, Alamo at 10 and 1 ppm, and Arbotect at 100, 10, and 1 ppm, were not significantly different in repression than the control ($P > 0.05$) (Table 2).

Experiment #1

Pathogenicity of fungal isolates

After inoculations, *L. longiclavatum* and *G. clavigera* successfully colonized the sapwood and resulted in mortality of 50 and 90% of seedlings, respectively (Table 1). One seedling in the DI water control group (10% mortality) and two in the sterile agar treatment group (20% mortality) died. As mentioned, these two control groups were pooled (15% mortality) for statistical comparisons, resulting in a significantly lower mortality than that observed with *G. clavigera* (FET, $P < 0.001$) but no difference in that observed with *L. longiclavatum* ($P = 0.467$) (Table 1). Ten percent mortality was observed in seedlings injected




Figure 1. Cross-sectional destructive autopsies of *P. monticola* seedlings conducted at 120 days: **A)** Comparison of a subset of blue-stain-fungi-inoculated seedlings (top, discolored) versus untreated controls (bottom, healthy). **B)** Water-inoculated seedling with healthy and intact vascular (xylem and phloem) tissues (40x). **C)** *Grosmannia clavigera*-inoculated seedling with discolored vascular tissues and necrotic phloem (40x). **D)** Clear, healthy sapwood from sterile agar-inoculated seedling (100x). **E)** *Leptographium longiclavatum*-inoculated seedlings with oxidized and discolored resin ducts and radial parenchyma (100x).

Table 2. *In vitro* screening of pesticides TREE-äge, Alamo, and Arbotect, measured as repression in mm distance from *G. clavigera* colony. Means and standard errors (SEs) are based on two treated discs in each of three dishes and result from an ANOVA with nested errors.

Treatment	Concentration (ppm)	mm suppression (\pm SE)
Control	0	4.3 \pm 2.8 cd ^a
TREE-äge	1	3.5 \pm 2.8 cd
TREE-äge	10	1.2 \pm 2.8 d
TREE-äge	100	0.3 \pm 2.8 d
TREE-äge	1000	1.2 \pm 2.8 d
TREE-äge	10000	3.7 \pm 2.8 cd
Alamo	1	0.7 \pm 2.8 d
Alamo	10	1.3 \pm 2.8 d
Alamo	100	20.2 \pm 2.8 b
Alamo	1000	32.8 \pm 2.8 a
Alamo	10000	38.8 \pm 2.8 a
Arbotect	1	4.0 \pm 2.8 cd
Arbotect	10	1.2 \pm 2.8 d
Arbotect	100	10.8 \pm 2.8 c
Arbotect	1000	21.0 \pm 2.8 b
Arbotect	10000	33.3 \pm 2.8 a

^a Treatment means with a letter in common are not significantly different based on the protected LSD procedure at significance level 0.05.

with Alamo and co-inoculated with *L. longiclavatum* and *G. clavigera*. This was compared to 50% and 90% in *L. longiclavatum* and *G. clavigera*, respectively, the reduction in mortality being significant when compared to that for seedlings inoculated with *G. clavigera* ($P = 0.007$) but not *L. longiclavatum* ($P > 0.05$). Observed percent foliage necrosis was 56.7 ± 14.5 and $91.0\% \pm 9.0$ SE in the *L. longiclavatum* and *G. clavigera*-inoculated seedlings compared to $20.8\% \pm 9.2$ SE in the Alamo/blue-stain inoculation (Table 3) and $20.5\% \pm 7.9$ SE in the pooled controls (DI water), respectively. Percent foliar necrosis was significantly higher for seedlings inoculated with *G. clavigera* than for Alamo/blue-stain-inoculated seedlings and the pooled controls (Table 3). Circumferential girdling was calculated by measuring lesion width in centimeters and dividing it by stem circumference. Lesions averaged 92.0 ± 5.7 , 51.8 ± 7.4 , 38.7 ± 5.7 , and $28.9\% \pm 6.4$ SE of stem

circumference for *G. clavigera*, *L. longiclavatum*, Alamo/blue-stain inoculation, and inoculated checks combined, respectively (Table 4). Girdling was significantly greater for seedlings inoculated with *G. clavigera* than for the other inoculation treatments and the pooled controls. Axial lesion lengths were greatest in *G. clavigera*-inoculated seedlings, followed by *L. longiclavatum*, Alamo/blue-stain, and controls (4.8 ± 0.5 , 2.3 ± 0.4 , 0.7 ± 0.1 , and $0.6 \text{ cm} \pm 0.1 \text{ SE}$, respectively) (Table 4). Dissections revealed that *G. clavigera* and *L. longiclavatum* inoculations had oxidized and brownish-blue discolored resin ducts, parenchyma ray cells, and phloem, while the sterile agar and water controls exhibited clear, light-colored sapwood and healthy phloem (Figure 1). The Alamo/blue-stain-inoculated seedlings also showed clear-light sapwood and healthy phloem, except for the dead seedlings, which exhibited dark vascular lesions matching those observed from *G. clavigera* and *L. longiclavatum*. Cross sections above and within the inoc-

Table 3. Percent foliar dieback in pathogenicity and phytotoxicity studies; means and standard errors (SEs) for each treatment are based on 10 seedlings, except for the inoculation controls (where the average of water and sterile agar treatment means and appropriate SE are reported).

Study	Treatment	Foliar dieback (%) Mean \pm SE
Pathogenicity	Alamo/B.S.	20.8 \pm 9.2 b ^z
	<i>G. clavigera</i>	91.0 \pm 9.0 a
	<i>L. longiclavatum</i>	56.7 \pm 14.5 ab
	Pooled controls	20.5 \pm 7.9 b
Phytotoxicity	TREE-äge	13.5 \pm 1.8 a
	Alamo	19.6 \pm 3.2 a
	Arbotect	25.0 \pm 4.2 a
	UTC	30.2 \pm 8.5 a

^z Treatments within a study with the same letter are not significantly different based on Tukey adjusted *P*-values.

ulation site of the controls indicated successful compartmentalization of incision wounds. *Grossmannia clavigera* and *L. longiclavatum*, identified by morphological analysis, were recovered from the excised cross sections.

Phytotoxicity of fungicides

After 120 days, mortality of seedlings treated with TREE-äge, Alamo, Arbotect, or untreated checks was 0%, 0%, 0%, and 10% mortality, respectively (Table 1), giving no indication of differences among pesticide treatments and control (FET, $P = 1.0$). Foliage necrosis did not differ significantly ($F_{3,36} = 1.98$, $P = 0.135$) among the untreated controls, TREE-äge, Alamo, and Arbotect treatments, with 30.2 ± 8.5 , 13.5 ± 1.8 , 19.6 ± 3.2 , and $25.0\% \pm 4.2 \text{ SE}$, respectively (Table 3). No foliar phytotoxicity was therefore observed due to application of any systemic chemistry at the rates used. Microscopic examination confirmed that discolored sapwood was present and limited in the TREE-äge, Alamo, and Arbotect treatments; resin ducts and phloem were neither oxidized nor necrotic. Cross sections of a single untreated control seedling that died had oxidized xylem and an unidentified infection, while all other untreated controls had clear sapwood and healthy phloem.

Experiment #2

Dual inoculation of *G. clavigera*

Inoculations of *G. clavigera* on both sides of seedling stems resulted in $91.0\% \pm 9.0 \text{ SE}$ seedling mortality, compared to 0.0% of the sterile agar controls. *G. clavigera* was identified by morphological characteristics and was re-isolated from cross sections of inoculated seedlings.

Table 4. Pathogenicity of inoculation treatments as measured by percent stem girdling and lesion length (cm).

	Treatment	N	Circumferential girdling (%) Mean \pm SE	Lesion length (cm) Mean \pm SE
Pathogenicity	Alamo/B.S.	10	38.7 \pm 5.7 b ^z	0.7 \pm 0.1 c ^{yz}
	<i>G. clavigera</i>	10	92.0 \pm 5.7 a	4.8 \pm 0.5 a
	<i>L. Longiclavatum</i>	6	51.8 \pm 7.4 b	2.3 \pm 0.4 b
	Pooled controls	8	28.9 \pm 6.4 b	0.6 \pm 0.1 c

^z Treatments with the same letter are not significantly different based on Tukey-adjusted *P*-values.

^y Mean separation results are based on analysis of log-transformed lesion heights.

DISCUSSION & CONCLUSIONS

The two Ophiostomatoid fungi isolated from western white pine, *Grosmannia clavigera* and *Leptographium longiclavatum*, caused mortality of *P. monticola* seedlings following inoculation. Seedlings faded as early as 40 days following inoculation with *G. clavigera*, whereas seedlings generally faded more slowly following inoculation with *L. longiclavatum*. These results may differ in mature trees, as trials were only performed on seedlings. However, Krokene and Solheim (1998) demonstrated that the results of inoculations of two- and four-year-old Norway spruce seedlings, with four bark beetle-associated blue-stain fungi, largely agreed with previous results of inoculated 40-year-old Norway spruce trees with the same fungal strains. These, as well as the current results, suggest that inoculation of seedlings can be a reliable bioassay to predict the pathogenicity of blue-stain fungi associated with bark beetles. As *L. longiclavatum* is a recently described species, information on its virulence is not extensive. Lee et al. (2006b) proved its pathogenicity on 98- to 130-year-old lodgepole pine (*P. contorta*), suggesting that it may contribute to the mortality of MPB-infested pines. Rice et al. (2007b) compared the virulence of the three main associated blue-stain fungi—*Ophiostoma montium*, *G. clavigera*, and *L. longiclavatum*—in lodgepole pine, jack pine, and lodgepole × jack hybrids, concluding that these species are about equally competitive and virulent, finding no significant differences in lesion lengths between *G. clavigera* and *L. longiclavatum*. The results of the current study differ from this, as a significant difference was observed in mortality and extent of circumferential lesions over time between the two species, with *G. clavigera* being more virulent. This difference could be due to the variability of virulence in geographically distant strains of *G. clavigera* and *L. longiclavatum*, attributed to host-specific effects (western white pine versus jack and lodgepole pines and their hybrids), or could also be attributed to the amount of inoculum used for infection. Further research including the addition of a second inoculation point on the same seedling with *L. longiclavatum*, may be helpful in elucidating this question.

This injection experiment used a newly developed technique, Micro I.V.'s, to inject the pesticide

treatments, and from researchers' observations it performed effectively. Of the 5 ml of TREE-äge, Alamo, and Arbotect placed into the Micro I.V.'s precise uptake into each seedling was not calculated. However, results indicate that the volume of solution administered and infused into the xylem tissues was sufficient to show effects in the xylem (e.g., lesion length). Seedling autopsies also revealed discoloration from dye, a component in the TREE-äge and Alamo formulations, visually confirming that the vascular tissues were exposed to the injected product. Seedling mortality was not observed following injection with TREE-äge, therefore, the method of injection and application of TREE-äge per se is not likely to cause injury to western white pines.

Since the isolates were obtained from TREE-äge-injected trees attacked by MPB, researchers strongly suspect the involvement of *G. clavigera* and *L. longiclavatum* and other MPB associates in tree mortality. Furthermore, it was demonstrated that *G. clavigera* and *L. longiclavatum* are pathogenic with differences in virulence on *P. monticola* seedlings. Field studies to evaluate tree protection using TREE-äge and Alamo in *P. monticola* trees attacked by MPB are currently underway, and will help determine the generality of these findings, and, more specifically, provide an indication of their relative importance in determining mortality of larger trees *in situ*.

In vitro studies provide useful information in the development of effective injection treatments for tree protection. These results demonstrate that TREE-äge alone did not provide any repression of *G. clavigera*, and are consistent with the recovery of isolates of *G. clavigera* and *L. longiclavatum* from treated trees. Repression of the fungi was evident in both Alamo and Arbotect injections, particularly at the higher concentrations (Table 2), suggesting that similar chemistries could be used for systemic treatment against blue-stain fungi. Research utilizing injection of both insecticide and fungicide chemistries is currently being conducted to evaluate protection against MPB/blue-stain associates *in situ* with *P. monticola*, the results of which will help to determine efficacy in tree protection. In the interim, it is advised that tree stewards and arborists consider both vector and microbial associates when treating trees with systemic pesticides.

Acknowledgments. This study was made possible through the support of the USDA FS, Southern Research Station, RWU-4552, Pineville, Louisiana, U.S.; Region 5 Forest Health Protection, Susanville, California, U.S.; and the Modoc National Forest, Alturas, California, U.S. The authors thank Cavell Brownie, Professor Emeritus, North Carolina State University, Raleigh North Carolina, U.S. for conducting the statistical analyses in this study. The authors also thank David Cox, Syngenta Crop Protection, LLC. for his review and comments. Any use of trade, product, or firm names is for descriptive purposes only and does not imply endorsement by the United States government.

LITERATURE CITED

- Adams, A.S., and D.L. Six. 2007. Temporal variation in mycophagy and prevalence of fungi associated with developmental stages of *Dendroctonus ponderosae* (Coleoptera: Curculionidae). *Environmental Entomology* 36:64–72.
- Amman, G.D. 1978. Biology, ecology, and causes of outbreaks of the mountain pine beetle in lodgepole pine forests. In theory and practice of mountain pine beetle management in lodgepole pine forests. pp. 39–53. In: D.L. Kibbee, A.A. Berryman, G.D. Amman, and R.W. Starks (Eds.). University of Idaho, Moscow, Idaho, U.S. Proceedings of a Symposium, 25–27 April 1978, Pullman, Washington, U.S.
- Amman, G.D., M.D. McGregor, and R.E. Dolph, Jr. 1989. Mountain pine beetle. USDA Forest Insect and Disease Leaflet 2. Washington, D.C.
- Basham, H.G. 1970. Wilt of loblolly pine inoculated with blue-stain fungi of the genus *Ceratocystis*. *Phytopathology* 60:750–754.
- Carroll, A.L., S.W. Taylor, J. Régnière, and L. Safranyik. 2003. Effects of climate change on range expansion by the mountain pine beetle in British Columbia. pp. 223–232. In: T.L. Shore, J.E. Brooks, and J.E. Stone (Eds.). Mountain Pine Beetle Symposium: Challenges and Solutions, Kelowna, BC, October 30–31, 2003. Can. For. Serv. Pac. For. Cent. Inf. Rep. BC-X-399.
- Doccola, J.J., B.L. Strom, C. Brownie, and K.D. Klepzig. 2011a. Impact of systemic fungicides on lesions formed by inoculation with the blue-stain fungus (*Ophiostoma minus*) in loblolly pine (*Pinus taeda* L.). *Arboriculture & Urban Forestry* 37(6):288–292.
- Doccola, J.J., D.R. Smitley, T.W. Davis, J.J. Aiken, and P.M. Wild. 2011b. Tree wound responses following systemic insecticide trunk injection treatments in green ash (*Fraxinus pennsylvanica* Marsh.) as determined by destructive autopsy. *Arboriculture & Urban Forestry* 37(1):6–12.
- Doccola, J.J., W. Hascher, J.J. Aiken, and P.M. Wild. 2012. Treatment strategies using hemlock woolly adelgid (*Adelges tsugae* Annand) infested eastern hemlock (*Tsuga canadensis* Carriere) trees. *Arboriculture & Urban Forestry* 38(2):41–49.
- Grosman, D.M., W.W. Upton, F.A. McCook, and R.F. Billings. 2002. Systemic insecticide injections for control of cone and seed insects in loblolly pine sees orchards—Two-year results. Texas Forest Service, Forest Pest Management, P.O. Box 310, Lufkin, Texas 75902–0310.
- Kim, J.J., E.A. Allen, L.M. Humble, and C. Breuil. 2005. Ophiostomatoid and basidiomycetous fungi associated with green, red, and grey lodgepole pines after mountain pine beetle infestation. *Canadian Journal of Forest Research* 35:274–284.
- Krokene, P., and H. Solheim. 1998. Assessing the virulence of four bark beetle-associated blue-stain fungi using Norway spruce saplings. *Plant Pathology* 47:537–540.
- Lee, S., J.J. Kim, and C. Breuil. 2005. *Leptographium longiclavatum* sp. nov., a new species associated with the mountain pine beetle, *Dendroctonus ponderosae*. *Mycology Research* 109:1162–1170.
- Lee, S., J.J. Kim, and C. Breuil. 2006a. Diversity of fungi associated with the mountain pine beetle, *Dendroctonus ponderosae*, and infested lodgepole pines in British Columbia. *Fungal Diversity* 22:91–105.
- Lee, S., J.J. Kim, and C. Breuil. 2006b. Pathogenicity of *Leptographium longiclavatum* associated with *Dendroctonus ponderosae* to *Pinus contorta*. *Canadian Journal of Forest Research* 36:2864–2872.
- Mathre, D.E. 1964. Pathogenicity of *Ceratocystis ips* and *Ceratocystis minor* to *Pinus ponderosa*. *Contrib. Boyce Thomson Inst. Plant Research* 22:363–388.
- McCullough, D.G., T.M. Poland, A.C. Anulewicz, P. Lewis, and D. Cappaert. 2011. Evaluation of *Agrilus planipennis* (Coleoptera: Buprestidae) control provided by emamectin benzoate and two neonicotinoid insecticides, one and two seasons after treatment. *Journal of Economic Entomology* 104(5):1599–1612.
- Ono, H. 2003. The mountain pine beetle: Scope of the problem and key issues in Alberta. pp. 62–66. In: T.L. Shore, J.E. Brooks, and J.E. Stone (Eds.). Mountain Pine Beetle Symposium: Challenges and Solutions, Kelowna, BC, 30–31 October 2003, Victoria, BC: Can. For. Serv. Pac. For. Cent. Inf. Rep. BC-X-399.
- Raffa, K.F., and A.A. Berryman. 1983. Physiological aspects of lodgepole pine wound response to a fungal symbiont of the mountain pine beetle *Dendroctonus ponderosae*. *The Canadian Entomologist* 115:723–734.
- Rice, A.V., and D.W. Langor. 2009. Mountain pine beetle-associated blue-stain fungi in lodgepole × jack pine hybrids near Grande Prairie, Alberta (Canada). *Forest Pathology* 39:323–334.
- Rice, A.V., M.N. Thormann, and D.W. Langor. 2007a. Mountain pine beetle associated blue-stain fungi cause lesions on jack pine, and lodgepole × jack hybrids in Alberta. *Canadian Journal of Botany* 85:307–315.
- Rice, A.V., M.N. Thormann, and D.W. Langor. 2007b. Virulence of, and interactions among, mountain pine beetle associated blue-stain fungi on two pine species and their hybrids in Alberta. *Canadian Journal of Botany* 46:1523–1527.
- Shigo, A.L. 1986. Trees and Microorganisms. A New Tree Biology: Facts, Photos, and Philosophies on Trees and Their Problems and Proper Care. Shigo and Trees, Associates, Durham, New Hampshire, U.S. pp.121–123.
- Shrimpton, D.M. 1973. Age- and size-related response of lodgepole pine to inoculation with *Europhium clavigerum*. *Canadian Journal of Botany* 51:1155–1160.
- Six, D.L., and T.D. Paine. 1998. Effects of mycangial fungi and host tree species on progeny survival and emergence of *Dendroctonus ponderosae* (Coleoptera: Scolytidae). *Environmental Entomology* 27:1393–1401.
- Six, D.L., T.C. Harrington, J. Steimel, D. McNew, and T.D. Paine. 2003. Genetic relationships among *Leptographium terebrantis* and the mycangial fungi of three western *Dendroctonus* bark beetles. *Mycologia* 95: 781–792.
- Smitley, D.R., J.J. Doccola, and D.L. Cox. 2010. Multiple-year protection of ash trees from emerald ash borer with a single trunk injection of emamectin benzoate, and single-year protection with an imidacloprid basal drench. *Arboriculture & Urban Forestry* 36(5):206–211.

- Solheim, H., and P. Krokene. 1998. Growth and virulence of mountain pine beetle associated blue-stain fungi, *Ophiostoma clavigerum* and *Ophiostoma montium*. Canadian Journal of Botany 76: 561–566.
- Strobel, G.A., and F. Sugawara. 1986. The pathogenicity of *Ceratocystis montia* to lodgepole pine. Canadian Journal of Botany 64:113–116.
- Tsuneda, A., and Y. Hiratsuka. 1984. Sympodial and annelidic conidiation in *Ceratocystis clavigera*. Canadian Journal of Botany 62:2618–2624.
- Wagener, W.W., and J.L. Mielke. 1961. A staining fungus root disease of ponderosa, Jeffrey, and pinyon pines. Plant Disease Reporter 45:831–835.
- Yamaoka, Y., R.H. Swanson, and Y. Hiratsuka. 1990. Inoculation of lodgepole pine with four blue-stain fungi associated with mountain pine beetle, monitored by a heat pulse velocity (HPV) instrument. Canadian Journal of Forest Research 20:31–36.
- Yamaoka, Y., Y. Hiratsuka, and P.J. Maruyama. 1995. The ability of *Ophiostoma clavigerum* to kill mature lodgepole pine trees. European Journal of Plant Pathology 25:401–404.

Stephen A. Wyka
Research Associate
Arborjet, Inc.
Woburn, Massachusetts 01801, U.S.

Joseph J. Doccia (corresponding author)
Director of Research & Development
Arborjet, Inc.
Woburn, Massachusetts 01801, U.S.
JoeDoccia@arborjet.com

Brian L. Strom
Research Entomologist
Southern Research Station
U.S. Department of Agriculture
Forest Service
2500 Shreveport Highway
Pineville, Louisiana 71360, U.S.

Sheri L. Smith
Regional Entomologist
R5 Forest Health Protection
U.S. Department of Agriculture
Forest Service
2550 Riverside Drive
Susanville, California 96130, U.S.

Douglas W. McPherson
Research Associate
Arborjet, Inc.
Woburn, Massachusetts 01801, U.S.

Srdan G. Acimović
Plant Pathologist
Arborjet, Inc.
Woburn, Massachusetts 01801, U.S.

Kier Klepzig
Assistant Director for Research
Southern Research Station, USFS
U.S. Department of Agriculture
Forest Service
200 WT Weaver Blvd.
Asheville, North Carolina 28804, U.S.

Résumé. Les scolytes transportent un certain nombre d'organismes qui se retrouvent sur l'arbre hôte lorsqu'ils l'attaquent et sont censés jouer un rôle dans le dépérissement des arbres. Afin d'évaluer la pathogénicité du pin argenté (*western white pine*, *Pinus monticola*) par les champignons transportés par le dendroctone du pin ponderosa (*mountain pine beetle*, *Dendroctonus ponderosae*), et afin d'évaluer le potentiel des traitements prophylactiques systémiques en vue de réduire les impacts des champignons, des expériences ont été menées sur des plantules de pin argenté afin de répondre à trois objectifs : 1) évaluer la pathogénicité de deux champignons du bleuissement associés au dendroctone du pin ponderosa; 2) évaluer la phytotoxicité de fongicides par injection dans les arbres; 3) et évaluer l'activité antifongique des produits injectés dans les arbres, *in vitro* et *in vivo*, à l'égard des champignons du bleuissement. Pour évaluer la pathogénicité, les plantules ont été inoculées avec le *Grosmannia clavigera* ou le *Leptographium clavatum*, maladies fongiques communes associées au dendroctone du pin ponderosa. La mortalité des semis quatre mois après l'inoculation était de 50% avec le *L. clavatum* et de 90% avec le *G. clavigera*, les deux significativement plus élevées que les plants témoins, démontrant ainsi la pathogénicité. Les effets phytotoxiques des fongicides *TREE-Age*®, *Alamo*® et *Arbotect*® furent évalués lors d'injections sur tige; aucun effet phytotoxique n'a été observé. Les propriétés anti-fongiques des trois mêmes produits ont été évaluées *in vitro* contre le *G. clavigera* alors que le fongicide *Alamo* s'est avéré le plus actif. La co-inoculation de *G. clavigera* et *L. longiclavatum* sur les plantules après injection sur tige du fongicide *Alamo* a démontrée significativement moins de mortalité et moins d'apparition de lésions que lorsqu'un seul des deux champignons est inoculé. Les résultats appuient l'hypothèse que les champignons du bleuissement associés au dendroctone du pin ponderosa, en particulier le *G. clavigera*, favorisent la mort de pin argenté quand il est attaqué par le dendroctone du pin ponderosa. Ces résultats suggèrent que l'application d'un fongicide en conjonction avec un insecticide pour la protection des arbres contre les scolytes peut être avantageuse.

Zusammenfassung. Borkenkäfer tragen eine Menge von assoziierten Organismen, die während der Attacke auf den Wirt übertragen werden und werden daher betrachtet, dass sie eine Rolle im Untergang der Bäume spielen. Um die Pathogenizität bei (WWP; *Pinus monticola*) ausgehend von Pilzregern, die von Bergkiefer-Käfern (MPB; *Dendroctonus ponderosae*) herumgetragen werden, zu untersuchen und das Potential für die Behandlung mit systemischen prophylaktischen Mitteln zur Reduzierung von Auswirkungen von Pilzkrankungen zu bewerten, wurden Experimente mit WWP-Sämlingen durchgeführt, die drei Ziele treffen sollen: 1. Bewertung der Pathogenizität von zwei mit MPB assoziierten Pilzen, die eine Verbläuung verursachen; 2. Bewertung der Phytotoxizität von drei Bauminjektions-Produkten; 3. Bewertung der Anti-Pilz-Aktivität von Bauminjektions-Produkten, *in vitro* und *in vivo*, gegen die assoziierten Bläue-Pilze. Um die Pathogenizität zu bewerten, wurden Sämlinge mit *Grosmannia clavigera* oder *Leptographium longiclavatum* inokuliert, zwei häufige Pilze in Zusammenhang mit den Bergkiefernkäfern (MPBs). Die Mortalität vier Monate nach der Inokulation betrug 50 % bei *L. longiclavatum* und

90 % bei with *G. clavigera*, beide signifikant höher als die Kontrollen und daher Pathogenizität demonstrierend. Die phytotoxischen Effekte von TREE-äge®, Alamo®, and Arbotect® wurden durch die Stamminjektion bewertet: es wurden keine phytotoxischen Effekte beobachtet. Von den selben drei Produkten wurden auch die Anti-Pilz-Aktivitäten in vitro gegen *G. clavigera* bewertet, wobei Alamo® die höchste Aktivität aufwies. Eine Co-Inoculation von *G. clavigera* und *L. longiclavatum* in Sämlinge nach einer Stamminjektion mit Alamo® zeigte signifikant weniger Mortalität und Verletzungsanzeichen als jede Spezies für sich allein. Die Resultate unterstützen die Hypothese, dass die Bläue-Pilzerreger, assoziiert mit den Bergkieferkäfern, insbesondere *G. clavigera*, ein Absterben von Bergkiefern herbeiführen, wenn diese durch den Käfer attackiert werden. Diese Ergebnisse lassen den Schluss zu, das die Handhabung eines Fungizids zusammen mit einem Insektizid für den Schutz von Bäumen gegen Borkenkäfer vorteilhaft wäre.

Resumen. Los insectos descortezadores transportan un sin número de organismos asociados que se transfieren al árbol huésped tras el ataque y se cree que desempeñan un papel en la declinación del árbol. Para evaluar la patogenidad para el pino blanco del oeste (WWP; *Pinus monticola*) de hongos transportados por el escarabajo del pino de montaña (MPB; *Dendroctonus ponderosae*) y para evaluar el potencial de los tratamientos profilácticos sistémicos para reducir los impactos de hongos, se realizaron experimentos con plántulas WWP para cumplir tres objetivos: 1) evaluar la patogenidad de dos hongos MPB de manchas azules asociados; 2) evaluar la fitotoxicidad de productos de inyección de árbol; 3) evaluar la actividad anti-fúngica de productos de inyección de árboles, in vitro y en vivo, hacia los hongos azules asociados. Para evaluar la patogenidad, las plántulas fueron inoculadas con *Grosmannia clavigera* o *Leptographium longiclavatum*, hongos comunes asociados de MPB. La mortalidad de las plántulas a los cuatro meses después de la inoculación fue de 50% con *L. longiclavatum* y 90% con *G. clavigera*, ambos significativamente más altos que los controles, demostrando así la patogenidad. Se evaluaron los efectos fitotóxicos de TREE-Age®, Alamo® y Arbotect® por inyección al tallo; no se observaron efectos fitotóxicos. Se evaluaron las propiedades anti-fúngicas de los mismos tres productos in vitro contra *G. clavigera* encontrando que Alamo era más activo. La co-inoculación de *G. clavigera* y *L. longiclavatum* en plántulas después de un inyección al tallo de Alamo mostraron significativamente menos mortalidad y la lesión de cualquiera de las especies. Los resultados apoyan la hipótesis de que la mancha azul asociada a MPB, particularmente *G. clavigera*, promueven la muerte de WWP cuando es atacado por MPB. Estos hallazgos sugieren que la administración de un fungicida con insecticida para la protección del árbol contra escarabajos de la corteza puede ser ventajoso.