



# Evaluation of Silicon Fertilizers and a Resistance Inducing Agent for Control of Apple and Pear Scab under Field Conditions

Glynn C. Percival

**Abstract.** Two field trials were conducted using established apple (*Malus* cv. Golden Delicious) and pear (*Pyrus communis* ‘Williams’ Bon Chrétien’) to assess the efficacy of a range of silicon (Si) fertilizers, a resistance-inducing (IR) agent based on salicylic acid and a combination of Si + IR when applied at four growth stages of tree development (i.e., bud break, green cluster, 90% petal fall, early fruitlet) against the foliar pathogens apple (*Venturia inaequalis*) and pear (*V. pirina*) scab. In addition, a comparative evaluation of a conventional synthetic fungicide (penconazole) used within the UK for apple and pear scab control was performed. Greatest protection against apple and pear scab was provided by a combination of a Si fertilizer + IR agent and the synthetic fungicide penconazole in both the 2014 and 2015 field trials where statistically comparable degrees of control were achieved. Efficacy as scab-protectant compounds (i.e., reduced scab severity of leaves and fruit) enhanced leaf chlorophyll content and fruit yield, and was confirmed when each Si fertilizer was applied at four growth stages during two growing seasons; however, there was little difference in the degree of control conferred by each Si fertilizer. The IR agent used alone proved the least effective scab protectant compound.

**Key Words.** Apple Scab; Fungicide; Integrated Disease Management; Orchard Management; Pathogen Control; Pear Scab; Plant Health Care; Urban Landscapes.

Apple and pear scab caused by the fungal pathogens *Venturia inaequalis* and *V. pirina* respectively can be devastating to apples (*Malus* spp.) and pears (*Pyrus* spp.), reducing fruit quality, marketable yield, and aesthetics (Cuthbertson and Murchie 2003; Jamar et al. 2010; Hailey and Percival 2014). The scab fungi infect leaves, petioles, and fruit, with initial symptoms appearing as olive-green spots on the leaf or leaf petiole. As infection develops, leaves turn yellow and drop prematurely, while infected fruit becomes deformed, scabby, and can drop before maturity (Jamar et al. 2010). Within Europe, suppliers, vendors, and growers of apples and pears adopt a zero-tolerance policy toward scab. As a result, the economics of fruit production require frequent application of synthetic fungicides throughout the growing season. Increased pathogen insensitivity to synthetic fungicides, coupled with public demands to reduce use, stimulated by greater awareness of environmental and health issues, has placed emphasis on the development of alternative control

strategies (Gozzo 2003; Villalta et al. 2004; Percival and Haynes 2008; Hailey and Percival 2014).

The potential benefits of silicon (Si) nutrition in plants include enhancement of growth and yield; improvement of mechanical properties (soil penetration by roots, resistance to lodging); reduced transpiration; and resistance to drought, salt, and heavy metal toxicities (Fauteux et al. 2005). In addition, Si-based soil fertilizers and sprays have proved effective in controlling soil-borne and foliar fungal pathogens of several plant species (Ma 2004). In rice, for example, Si has been demonstrated to control rice blast, (*Magnaporthe grisea*, anamorph = *Pyricularia grisea*), as effectively as commercially available fungicides as well as reduce the number of fungicide applications (Datnoff et al. 2001; Seebold et al. 2004). Studies of the influence of Si fertilizers on gray leaf spot development in St Augustine grass under greenhouse conditions demonstrated that Si significantly reduced disease progress by 44% to 78% and whole-plant infection by 2.5% to 50.5%.

(Datnoff and Nagata 1999). Si fertilizers also have been shown to reduce the incidence of powdery mildew in Kentucky bluegrass (Hamel and Heckman 2000), improve disease resistance in zoysia grass to *Rhizoctonia solani* (Saigusa et al. 2000) and creeping bentgrass to *Pythium aphanidermatum*, *Sclerotinia homoeocarpa* and *R. solani* (Schmidt et al. 1999; Rondeau 2001; Uriarte et al. 2004).

Induced resistance (IR) is the phenomenon whereby a plant's own defense mechanisms are induced by treatment with a biological (i.e., weakened or attenuated fungal pathogen) or chemical agent (i.e., inorganic potassium and phosphate salts, low-molecular-weight proteins, unsaturated fatty acids; (Bécot 2000; Fobert and Després 2005; Percival et al. 2009). Developments in plant protection technology have led to the formulation of a range of commercially available IR compounds such as harpin protein, potassium phosphite, and a range of salicylic acid analogs that are registered for commercial use within the horticultural industry, although their availability differs between countries (Percival and Haynes 2008). Studies show that IR compounds can reduce disease severity of fungal pathogens, such as *Venturia inaequalis* and *V. pirina*, with the level of pathogen suppression, on occasion, comparable with synthetic fungicides (Christiansen et al. 1999; Bernards and Bastrup-Spohr 2008; Percival et al. 2009). The potential of Si fertilizers singly and in combination with an inducing agent to manage scab diseases of trees has received little study. The majority of studies investigating the efficacy of IR agents on pathogen suppression were conducted under controlled laboratory and glasshouse conditions that do not reflect field environments (Agnostini et al. 2003; Percival et al. 2009). The purpose of this study was to investigate the association of Si fertilizers singly and in combination with an IR agent as scab-protective compounds, using scab-sensitive species of apple (*Malus* cv. Golden Delicious) and pear (*Pyrus communis* 'Williams' Bon Chrétien').

## MATERIALS AND METHODS

### Field Trials

The apple trial site consisted of a 0.75 ha block of apple (*Malus* cv. Golden Delicious) interspersed with individual trees of *Malus* Red Delicious and

Gala as pollinators. The pear trial site consisted of a 0.90 ha block of *Pyrus communis* 'Williams' Bon Chrétien' interspersed with individual trees of *Pyrus communis* Beth and Concorde. Planting distances were 2 m × 2 m spacing. Trees were planted in 2003 and trained under a central-leader system to an average height of 2.5 m ± 0.25 m with mean trunk diameters of 12 cm ± 1.4 cm at 45 cm above the soil level. The trial sites were located at the University of Reading Shinfield Experimental Site, Berkshire, UK (51°43N, -1°08W).

The soil was a sandy loam containing 4%–6% organic matter; pH of 6.2; and available P, K, Mg, Na, and Ca were 52.0, 659.1, 175.2, 49.4, and 2,188 mg l<sup>-1</sup>, respectively. Weeds were controlled using glyphosate (Roundup®; Green-Tech, Sweethills Park, Nun Monkton, York, UK). No watering or fertilization was applied during the two-year trial. *Venturia inaequalis* and *V. pirina* occur every year due to monocultures in the experimental area where the experiments were carried out. Prior to studies commencing in 2014 and 2015, trees were inspected in September 2013 and 2014, and only those trees with >50% of leaves affected with severe foliar discoloration and scab infection were used. A minimal insecticide program, based on deltamethrin (Bandu, Headland Agrochemicals Ltd, Saffron Walden, Essex, UK), was applied every two months from May 2014 to September 2015. All sprays were applied using a Tom Wanner Spray Rig sprayer at 40 ml deltamethrin per 100 l<sup>-1</sup> of water. Trees were sprayed until runoff, 0.30 l<sup>-1</sup> insecticide per tree.

### Silicon and IR Treatments

Treatments were applied at four distinct growth stages (Bevan and Knight 2001), namely: bud break (07 March 2014, 18 March 2015), green cluster (01 April 2014, 06 April 2015), 90% petal fall (08 May 2014, 21 May 2015), and early fruitlet (03 June 2014, 14 June 2015). Prior to each treatment, trees were inspected, and no symptoms of apple or pear scab were apparent. During spray treatments, polythene screens 2.5 m high were erected around each tree to prevent spray drift and cross contact with other trees. The base of the tree was covered with a 0.5 m × 0.5 m polythene mulch to prevent potential soil percolation. The treatments, 1 IR compound, 1 fungicide, 4 silicon fertilizers × 4 spray times were applied in 10

randomized complete blocks plus a water control with a single tree as the experimental unit, giving a total of 60 observations per response variable. Foliar sprays of each Si fertilizer, IR agent, and penconazole were applied until runoff using a 10 liter knapsack sprayer (Cooper Pedlar, Agratech NW Ltd, Waterfoot, Rossendale, UK) at:

- Topas (a.i. penconazole): 1.5 ml l<sup>-1</sup> of water (Syngenta Crop Protection UK Ltd, Whittlesford, Cambridge, UK).
- Rigel (a.i. Polyether modified trisiloxane + Salicylaldehyde): 3 ml l<sup>-1</sup> of water (Orion Future Technology Ltd, Henwood House, Henwood, Ashford, Kent, UK).
  - ▶ Equivalent to 7 g Si per liter of water
- Sirius (tetraethyl orthosilicate [Si (OC<sub>2</sub>H<sub>5</sub>)<sub>4</sub>]): 10 ml l<sup>-1</sup> of water (Orion Future Technology Ltd, Henwood House, Henwood, Ashford, Kent, UK).
  - ▶ Equivalent to 7 g Si per liter of water
- Fossil (polyether modified trisiloxane): 10 ml l<sup>-1</sup> of water (Orion Future Technology Ltd, Henwood House, Henwood, Ashford, Kent, UK).
  - ▶ Equivalent to 7 g Si per liter of water
- Salicylaldehyde (Sigma-Aldrich, Gillingham, Dorset, UK).
- Sil-MATRIX (potassium silicate) 10 ml l<sup>-1</sup> of water (Certis Granta Park, Riverside, Great Abington, Cambridge, UK).

## Tree Vitality

### Leaf chlorophyll measurements

To keep the age of the leaves comparable throughout the experiment, measurements of chlorophyll content (SPAD) were made only on fully expanded mature leaves. In all cases measurements were taken from six leaves (two from the top of the crown, two in the center, and two at the base) per tree. A chlorophyll meter was used (SPAD-502 Chlorophyll Meter, Konica Minolta Sensing Europe B.V.). Chlorophyll was measured at the midpoint of the leaf next to the main leaf vein. Calibration was obtained by measurement of absorbance at 663 and 645 nm in a spectrophotometer (PU8800 Pye Unicam) after extraction with 80% v/v aqueous acetone (regr. eq. = 5.75 + 0.055x; r<sup>2</sup> adj = 0.96, P ≤ 0.001) (Lichtenthaler and Wellburn 1983).

### Scab severity

Scab severity of leaves and fruit was assessed 28 September 2014 and 28 September 2015. Leaf scab severity of each tree was rated using a visual indexing technique and ratings on the scale: 0 = No scab observed; 1 = less than 5% of leaves affected and no aesthetic impact; 2 = 5%–20% of leaves affected with some yellowing but little or no defoliation; 3 = 21%–50% of leaves affected, significant defoliation and/or leaf yellowing; 4 = 51%–80% of leaves affected, severe foliar discoloration; 5 = 81%–100% of leaves affected with 90%–100% defoliation. Scab severity on fruit was calculated on the scale: 0 = no visible lesions; 1 = <10% fruit surface infected; 2 = 10%–25% fruit surface infected; 3 = 25%–50% fruit surface infected; 4 = >50% fruit surface infected. The individual ratings for each tree in each treatment were used as a measure of scab severity for statistical analysis. Leaf scab severity ratings used in this study was based on UK and Ireland market standards for fungicide evaluation of scab control (Butt et al. 1990; Swait and Butt 1990). Fruit scab severity was based on a scale used by Ilhan et al. (2006).

### Fruit Yield

Yield per tree was determined by weighing all fruit on each tree at harvest and dividing by the number of trees per treatment.

### Silicon Analysis

Ten leaves and three fruits per tree were collected for tissue Si analysis. The leaves and fruit per tree were pooled for randomization purposes, washed with deionized water, and then placed in an oven for three days at 70°C. Dried leaves and fruit were then ground to pass through a 40-mesh screen. Leaf tissue Si concentration was determined following the method of Elliott and Snyder (1991). Leaves were ashed on platinum dishes and ash mixed with Na<sub>2</sub>CO<sub>3</sub> and H<sub>3</sub>BO<sub>3</sub> (5:1) to melt. Aliquot product was dissolved in HCl solution and transferred to a volumetric flask. Ammonium molybdate, tartaric acid, and ascorbic acid were added to this solution, and the absorbance of the solution measured at 420 and 655 nm using a spectrophotometer (UV-1600; SHIMAZDU Co., Kyoto, Japan). The silica content was calculated per unit area and on a dry weight basis by comparing the absorbance with standard silicon solutions (10–1000 ppm).

## Statistical Analysis

Mean pathogen severity values for all treatments were transformed using the Arcsin (sine<sup>-1</sup>) transformation. All data were analyzed using ANOVA, and the differences between means were determined using Tukey *w* procedure ( $P = 0.05$ ), Genstat® for Windows® program. Back-transformed pathogen severity values are presented here to ease interpretation of data (Blaedow et al. 2006).

## RESULTS

Damaging outbreaks of apple and pear scab were recorded on foliage of control trees in both the 2014 and 2015 trials as indicated by leaf scab severity ratings of 4.8 and 5.0 on *Malus* cv. Golden Delicious, and 3.4 and 3.4 on *Pyrus communis* 'Williams' Bon Chrétien' at the end of the 2014 and 2015 growing season, respectively (Table 1; Table 2). None of the treated

or control trees died as a result of scab attack during the course of the two-year study (data not shown).

Efficacy as scab-protectant compounds (i.e., reduced scab severity of leaves and fruit) enhanced leaf chlorophyll content and fruit yield was confirmed when penconazole (synthetic fungicide), salicylaldehyde (IR agent), Sil-MATRIX, Fossil, Sirius (Si fertilizers), and Rigel (Si fertilizer + IR agent) were applied at four growth stages during two growing seasons (Table 1; Table 2; Table 3; Table 4). In most instances, these effects were statistically significant ( $P < 0.05$ ; Table 1; Table 2; Table 3; Table 4). Maximal reductions in apple and pear scab severity on leaves and fruit was recorded following application of the synthetic fungicide penconazole and a combination of a polyether modified trisiloxane (Si fertilizer) + salicylaldehyde (IR agent; trade name Rigel) (Tables 1; Table 2). In all cases there was no

**Table 1. Selected silicon fertilizers and inducing agents applied by foliar sprays compared for the control of *Venturia inaequalis* and *V. pirina* on leaves of apple (*Malus* cv. Golden Delicious) and pear (*Pyrus communis* 'Williams' Bon Chrétien'), respectively, as measured by an observed leaf scab severity index<sup>z</sup>.**

Treatment	Apple scab severity on leaves <sup>y</sup>		Pear scab severity on leaves <sup>y</sup>	
	2014	2015	2014	2015
Water (control)	4.8a	5.0a	3.4a	3.4a
Sil-MATRIX (10 ml liter of water)	2.7bc	3.0bc	2.4b	2.4b
Salicylaldehyde	3.0b	3.2b	2.2b	2.4b
Fossil (10 ml liter of water)	2.3bc	2.2bcd	1.9b	2.0bc
Sirius (10 ml liter of water)	2.3bc	2.4bcd	1.8b	2.0bc
Rigel (3 ml liter of water)	1.6cd	1.8de	1.6bc	1.4cd
Topas (1.5 ml liter of water)	0.4d	0.8e	0.8c	1.0d
Treatment	<0.001	<0.001	<0.001	<0.001

<sup>z</sup> Scab severity index: 0 = no scab observed; 1 = less than 5% of leaves affected and no aesthetic impact; 2 = 5%–20% of leaves affected with some yellowing but little or no defoliation; 3 = 21%–50% of leaves affected, significant defoliation, and/or leaf yellowing; 4 = 51%–80% of leaves affected, severe foliar discoloration; 5 = 81%–100% of foliage affected with 90%–100% defoliation.

<sup>y</sup> Mean of ten randomized complete blocks with one tree per block.

Note: Lowercase letters indicate significant differences between means for each evaluation date by Tukey highly significance test ( $P = 0.05$ ).

**Table 2. Selected silicon fertilizers and inducing agents applied by foliar sprays compared for the control of *Venturia inaequalis* and *V. pirina* on fruit of apple (*Malus* cv. Golden Delicious) and pear (*Pyrus communis* 'Williams' Bon Chrétien'), respectively, as measured by an observed scab severity index<sup>z</sup>.**

Treatment	Apple scab severity on fruit <sup>y</sup>		Pear scab severity on fruit <sup>y</sup>	
	2014	2015	2014	2015
Water (control)	2.3a	2.5a	2.1a	1.8a
Sil-MATRIX (10 ml liter of water)	1.9ab	1.8ab	1.5b	1.3b
Salicylaldehyde	1.4bc	1.6b	1.4bc	1.4ab
Fossil (10 ml liter of water)	1.5bc	1.2bc	1.2bc	1.0bc
Sirius (10 ml liter of water)	1.2cd	1.5bc	0.9cd	1.1bc
Rigel (3 ml liter of water)	0.7de	0.9cd	0.6d	0.7c
Topas (1.5 ml liter of water)	0.2e	0.4d	0.4d	1.0c
Treatment	<0.001	<0.001	<0.001	0.001

<sup>z</sup> Scab severity index: 0 = no scab observed; 1 = less than 5% of leaves affected and no aesthetic impact; 2 = 5%–20% of leaves affected with some yellowing but little or no defoliation; 3 = 21%–50% of leaves affected, significant defoliation, and/or leaf yellowing; 4 = 51%–80% of leaves affected, severe foliar discoloration; 5 = 81%–100% of foliage affected with 90%–100% defoliation.

<sup>y</sup> Mean of ten randomized complete blocks with one tree per block.

Note: Lowercase letters indicate significant differences between means for each evaluation date by Tukey highly significance test ( $P = 0.05$ ).

statistically significant ( $P < 0.05$ ) differences between these two treatments. The mean over two growing seasons reductions in leaf and fruit scab severity ranged from 62.5%–91.6% and 60.8%–91.3% (*Malus* cv. Golden Delicious) and from 52.9%–76.4% and 52.4%–81.0% (*P. communis* ‘Williams’ Bon Chrétien’) respectively (Table 1; Table 2). The mean over two growing seasons increased in leaf chlorophyll content (SPAD values) and fruit yield per tree (kg), and ranged from 47.3%–64.1% and 46.1%–65.4% (*Malus* cv. Golden Delicious) and from 36.6%–53.4% and 40.0%–53.3% (*P. communis* ‘Williams’ Bon Chrétien’), respectively (Table 3; Table 4).

Applications of each Si fertilizer namely; tetraethyl orthosilicate [ $\text{Si}(\text{OC}_2\text{H}_5)_4$ ] (trade name Sirius), polyether modified trisiloxane (trade name Fossil), potassium silicate (trade name Sil-MATRIX) reduced leaf and fruit scab severity by 37.5%–54.1% and 20.8%–50.0% (*Malus* cv. Golden Delicious) and by 29.4%–47.1% and 23.1%–53.8% (*Pyrus communis* ‘Williams’ Bon Chrétien’), respectively, when averaged over both growing seasons

(Table 1; Table 2). Improvements in leaf chlorophyll content SPAD values ranged from 15.4%–49.2% (*Malus* cv. Golden Delicious) and 9.0%–21.0% (*Pyrus communis* ‘Williams’ Bon Chrétien’) (Table 3), while improvements in *Malus* cv. Golden Delicious and *Pyrus communis* ‘Williams’ Bon Chrétien’ fruit yield (kg per tree) ranged from 5.5%–27.3% and 17.3%–34.5%, respectively, compared to water-treated control trees when averaged over two growing seasons (Table 4). Of all the products evaluated, the IR agent salicylaldehyde proved the least effective scab-protectant compound reducing leaf and fruit scab severity by 36.7% and 37.5% (*Malus* cv. Golden Delicious) and 32.4% and 28.2% (*Pyrus communis* ‘Williams’ Bon Chrétien’), respectively, when averaged over two growing seasons (Table 1; Table 2). Improvements in leaf chlorophyll content SPAD values and fruit yield (kg per tree) were, however, comparable with all Si treatments with values of 23.1% (*Malus* cv. Golden Delicious) and 12.9% (*Pyrus communis* ‘Williams’ Bon Chrétien’) and 10.9% and 18.1%, respectively, recorded com-

**Table 3. Selected silicon fertilizers and inducing agents applied by foliar sprays compared for the control of *Venturia inaequalis* and *V. pirina* on fruit of apple (*Malus* cv. Golden Delicious) and pear (*Pyrus communis* ‘Williams’ Bon Chrétien’), respectively, as measured by leaf chlorophyll content SPAD values.**

Treatment	Apple scab SPAD <sup>z</sup>		Pear scab SPAD <sup>z</sup>	
	2014	2015	2014	2015
Water (control)	24.5a	27.3a	29.2a	32.2a
Sil-MATRIX (10ml liter of water)	30.0ab	32.4ab	33.8ab	37.5ab
Salicylaldehyde	31.8b	33.2abc	34.1abc	36.4ab
Fossil (10 ml liter of water)	34.0b	37.1bcd	35.5abc	34.9a
Sirius (10 ml liter of water)	31.2a	38.8bcd	36.9bc	37.2ab
Rigel (3 ml liter of water)	36.1b	39.4cd	39.9bc	42.6bc
Topas (1.5 ml liter of water)	36.4b	40.2d	41.4c	44.8c
Treatment	0.035	0.044	0.031	0.050

<sup>z</sup> Mean of ten randomized complete blocks with one tree per block. Ten leaves per tree were randomly selected throughout the crown for chlorophyll content measurements, and the mean was calculated per tree for statistical purposes.

Note: Lowercase letters indicate significant differences between means for each evaluation date by Tukey highly significance test ( $P = 0.05$ ).

**Table 4. Selected silicon fertilizers and inducing agents applied by foliar sprays compared for the control of *Venturia inaequalis* and *V. pirina* on mean fruit yield of apple (*Malus* cv. Golden Delicious) and pear (*Pyrus communis* ‘Williams’ Bon Chrétien’), respectively.**

Treatment	Apple scab yield/tree (kg) <sup>z</sup>		Pear scab yield/tree (kg) <sup>z</sup>	
	2014	2015	2014	2015
Water (control)	5.2a	5.9a	6.0a	5.6a
Sil-MATRIX (10 ml liter of water)	5.8ab	7.0ab	7.2ab	6.8ab
Salicylaldehyde	6.0ab	6.2a	6.8a	6.9ab
Fossil (10 ml liter of water)	5.8ab	6.6ab	7.4abc	7.4bc
Sirius (10 ml liter of water)	6.4b	6.7ab	7.3ab	7.8bcd
Rigel (3 ml liter of water)	7.9c	7.6b	8.8c	9.2d
Topas (1.5 ml liter of water)	8.6c	7.8b	8.4c	8.4cd
Treatment	0.030	0.045	0.050	<0.022

<sup>z</sup> Mean of ten randomized complete blocks with one tree per block.

Note: Lowercase letters indicate significant differences between means for each evaluation date by Tukey highly significance test ( $P = 0.05$ ).

pared to water-treated control trees when averaged over two growing seasons (Table 3; Table 4).

In all cases, application of a Si fertilizer (i.e., Sil-MATRIX, Fossil, Sirius, Rigel) significantly ( $P < 0.05$ ) increased leaf and fruit Si content in both *Malus* cv. Golden Delicious and *Pyrus communis* 'Williams' Bon Chrétien' compared to water-treated controls. However, there was no major difference in the level of Si content recorded between each Si fertilizer (Table 5; Table 6). Within leaf tissue of *Malus* cv. Golden Delicious and *Pyrus communis* 'Williams' Bon Chrétien', Si content increased by 36.2%–58.5% and 34.5%–53.4%, respectively (Table 5). Within fruit tissue of *Malus* cv. Golden Delicious and *Pyrus communis* 'Williams' Bon Chrétien', Si content increased by 57.0%–104.2% and 76.9%–107.4%, respectively (Table 6). Application of penconazole or salicylaldehyde had no significant effect on leaf and fruit Si content of both *Malus* cv. Golden Delicious and *Pyrus communis* 'Williams' Bon Chrétien' where values were statistically comparable to controls. (Table 5; Table 6).

## DISCUSSION

A significant body of literature exists that describes Si nutrition as reducing the intensity of fungal diseases in peach, grape, barley, corn, rice, rye, strawberry, and wheat (Datnoff et al. 2007; Resende et al. 2012). However, few studies have investigated increased host resistance to scab epidemics promoted by Si under field conditions. Some authors suggest that mechanical or physical barriers promoted by Si deposition in cell walls contributes to enhanced resistance (Kim et al. 2002; Hayasaka et al. 2008). For example, Si application led to a pronounced cell silicification in rice leaves and more elaborate and larger papillae, with the authors concluding that the elaborate papillae formed in Si-treated leaf epidermal surface might increase host resistance to fungal penetration (Zhang et al. 2006; Cai et al. 2008). In addition, enhanced Si layers were observed in Si-treated plant epidermal cell walls, increasing wall thickness and conferring enhanced host resistance. Further studies have shown enhanced cuticular Si layer development on plants supple-

**Table 5. The influence of silicon fertilizers and inducing agents applied by foliar sprays on silicon content (mg/kg) of leaves of apple (*Malus* cv. Golden Delicious) and pear (*Pyrus communis* 'Williams' Bon Chrétien').**

Treatment	Apple <sup>z</sup>		Pear <sup>z</sup>	
	2014	2015	2014	2015
Water (control)	22.8a	26.4a	26.2a	20.1a
Sil-MATRIX (10 ml liter of water)	35.3b	38.1b	34.8b	31.2b
Salicylaldehyde	23.7a	23.9a	24.2a	18.7a
Fossil (10 ml liter of water)	36.0b	35.2b	35.0b	32.8bc
Sirius (10 ml liter of water)	36.5b	39.0b	35.6b	34.6bc
Rigel (3 ml liter of water)	33.5b	37.2b	33.3b	35.5c
Topas (1.5 ml liter of water)	20.1a	24.1a	22.8a	19.5a
Treatment	0.001	0.001	0.001	0.001

<sup>z</sup> Mean of ten randomized complete blocks with one tree per block. Ten leaves per tree were randomly selected throughout the crown for Si analysis. Note: Lowercase letters indicate significant differences between means for each evaluation date by Tukey highly significance test ( $P = 0.05$ ).

**Table 6. The influence of silicon fertilizers and inducing agents applied by foliar sprays on silicon content (mg/kg) of fruit of apple (*Malus* cv. Golden Delicious) and pear (*Pyrus communis* 'Williams' Bon Chrétien').**

Treatment	Apple <sup>z</sup>		Pear <sup>z</sup>	
	2014	2015	2014	2015
Water (control)	15.0a	13.4a	14.6a	12.3a
Sil-MATRIX (10 ml liter of water)	26.1bc	24.8b	27.9b	23.8b
Salicylaldehyde	16.8a	15.3a	13.9a	14.6a
Fossil (10 ml liter of water)	22.3b	27.0b	25.0b	25.0bc
Sirius (10 ml liter of water)	26.5bc	25.8b	24.3b	27.1bc
Rigel (3 ml liter of water)	29.0c	26.1b	27.2b	27.5c
Topas (1.5 ml liter of water)	17.4a	12.4a	14.0a	14.1a
Treatment	0.001	0.001	0.001	0.001

<sup>z</sup> Mean of ten randomized complete blocks with one tree per block. Three fruits per tree were randomly selected throughout the crown for Si analysis. Note: Lowercase letters indicate significant differences between means for each evaluation date by Tukey highly significance test ( $P = 0.05$ ).

mented with Si fertilizers constituted a physical barrier to impede fungal penetration and colonization (Kim et al. 2002; Hayasaka et al. 2008). Other studies, suggest that Si plays a biochemical role (i.e., Si-induced accumulation of phenolic compounds, phytoalexins, lignin, and peroxidase) in mediating plant resistance to pathogens (Rodrigues et al. 2003; Liang et al. 2005; Sun et al. 2010). In the current study, a significant increase in leaf and fruit Si content was recorded in both *Malus* cv. Golden Delicious and *Pyrus communis* 'Williams' Bon Chrétien' following application of any of the Si fertilizers. Such a response is consistent with other findings following application of Si fertilizers to peach, grape, kiwi, as well as several economically important grain and grass crops (Catherine Keller and Meunier 2012). Enhanced Si content in leaves and fruit suggests that Si-enhanced scab resistance may be partially caused by the role of Si as a physical barrier (Sun et al. 2010), coupled with promotion of inherent biochemical plant defense systems in apple and pear not investigated here.

Penconazole applied at bud break, green cluster, petal fall, and early fruitlet formation was highly effective in reducing apple and pear scab severity of fruit and leaves. The effectiveness of penconazole against apple and pear scab under laboratory and field conditions has been confirmed elsewhere (Schnabel and Parisi 1997; Percival et al. 2009). However, due to the development of triazole-insensitive strains of scab, greater emphasis on alternative scab management strategies is now warranted (Gozzo 2003; Vallad and Goodman 2004; Fobert and Després 2005; Witzell and Martin 2008). The phenomenon of inducing resistance in plants by biological, natural, and/or inorganic compounds potentially offers an alternate approach to scab management (Hammerschmidt 2003; Percival and Haynes 2008; Percival et al. 2009; Hailey and Percival 2014). Consequently, developments in plant protection technology have focused on the production of commercially available IR agents. In this study, the IR agent salicylaldehyde reduced leaf and fruit scab severity by ca. 37.0% (*Malus* cv. Golden Delicious) and 30% (*Pyrus communis* 'Williams' Bon Chrétien'). Salicylic acid (SA) and SA derivatives are phenolic-based com-

pounds found in a wide range of plant species that directly or indirectly influence plant metabolism. Application of SA to plants induces synthesis of low-molecular-weight (LMW) compounds (phenolics, terpenoids, alkaloids) that possess toxic, antimicrobial, anti-nutritive, and anti-digestive activity (Facchini 2001; Vallad and Goodman 2004; Fauteux et al. 2005; Keeling and Bohlmann 2006; Bernards and Bastrup-Spohr 2008; Witzell and Martin 2008). Consequently, stimulation and elevated concentrations of these compounds within plant tissue can significantly contribute to reduced severity of fungal, bacterial, and viral pathogens (Bokshi et al. 2003; Sparla et al. 2004).

Results of this study demonstrate that the commercially available Si fertilizer Rigel (a.i. polyether modified trisiloxane + salicylaldehyde) was as effective a scab-protectant product as penconazole. In all cases, reductions in scab leaf and fruit severity and enhancement of leaf color and fruit yield were statistically comparable to penconazole when applied four times from bud break to early fruitlet formation. In comparison to all other Si fertilizers evaluated in this study and the IR agent salicylaldehyde alone, a combination of Si + IR proved superior to Si or IR applied individually. Consequently, this indicates additive and/or synergistic effects when combining an Si fertilizer with an IR agent. From a commercial aspect, producers, suppliers, and vendors of apples generally adopt a low-tolerance policy toward apple and pear scab on fruit (Butt et al. 1990). Combining a Si + IR agent reduced scab levels to those achieved using penconazole, a synthetic fungicide. Such a result would prove of interest to professionals involved in the production, supply, and/or consumption of apple and pear especially where fruit produce is sold under an organic or naturally produced label, as scab severity levels tend to be less stringent (Bevan and Knight 2001) for ornamental apples planted for aesthetic reasons within town and city landscape industry, lower scab levels are also acceptable (Percival et al. 2009). Because of their mode of action, aimed at enhancing the physical and biochemical defense mechanisms of treated host plants rather than at directly arresting or killing a fungal disease agent, an Si fertilizer + IR agent potentially offers opportunities for the control of

other fungal pathogens in natural ecosystems. As effects are mostly on the plant itself, unlike with a synthetic fungicide, little, if any, consequences on the existing tree microbial community exist. In addition, Si fertilizers and/or IR agents are potential candidates for treatment in natural ecosystems because they have extremely low toxicity to invertebrates, aquatic organisms, or animals, including humans, compared to synthetic fungicides (Vallad and Goodman 2004; Garbelotto et al. 2007). Finally, Si fertilizers and/or IR agents can be incorporated into an integrated pest management system and/or be used preventatively to bolster general plant health (Sparla et al. 2004; Walters 2009; Percival 2016). In these instances, the reductions in scab severity recorded in this investigation may warrant the use of an Si fertilizer and/or IR agents as an alternative or complement to conventional synthetic fungicides.

In conclusion, the results of this study provide some evidence that Si fertilizers and/or IR agents individually or in combination could potentially play a useful role as an alternative and/or supplementary method of apple and pear scab management under field and/or landscape conditions, provided four sprays are applied from bud break to early fruitlet formation.

**Acknowledgments.** The author is grateful for funding in part from TREE FUND (Hyland Johns Grant).



## LITERATURE CITED

- Agostini, J.P., P.M. Bushong, and L.W. Timmer. 2003. Glasshouse evaluation of products that induce host resistance for control of scab, melanose, and *Alternaria* brown spot of citrus. *Plant Disease* 87:69–75.
- Bécot, S., E. Pajot, D. Le Corre, C. Monot, and D. Silué. 2000. Phytogard ( $K_2HPO_3$ ) induces localized resistance in cauliflower to downy mildew of crucifers. *Crop Protection* 19:417–425.
- Bernards, M.A., and L. Bastrup-Spohr. 2008. Phenylpropanoid metabolism induced by wounding and insect herbivory. pp. 189–213 In: A. Schaller (Ed.). *Induced plant resistance to herbivory*. Springer, New York, New York, U.S. 462 pp.
- Bevan, J., and S. Knight. 2001. *Organic Apple Production*, first edition. HDRA, Emmerson Press, Kenilworth, UK.
- Blaedow, R.A., W.R. Chaney, P.C. Pecknold, and H.A. Holt. 2006. Investigation of fungicidal properties of the tree growth regulator paclobutrazol to control apple scab. *Journal of Arboriculture* 32:67–73.
- Butt, D.J., A.A.J. Swait, and J.D. Robinson. 1990. Evaluation of fungicides against apple powdery mildew and scab. *Tests of Agrochemicals and Cultivars* 11. *Annals Applied Biology* (supp) 116:34–35.
- Cai, K.Z., D. Gao, S.M. Luo, R.S. Zeng, and J.Y. Yang. 2008. Physiological and cytological mechanisms of silicon-induced resistance in rice against blast disease. *Physiologia Plantarum* 134:324–333.
- Catherine Keller, F.G., and J.D. Meunier. 2012. Benefits of plant silicon for crops: A review. *Agronomy Sustainable Development* 32:201–213.
- Christiansen, E., P. Karokene, A.A. Berryman, V.R. Franceschi, T. Krekling, F. Lieutier, and A. Lonneborg, et al. 1999. Mechanical injury and fungal infection induce acquired resistance in Norway spruce. *Tree Physiology* 19:399–403.
- Cuthbertson, A.G.S., and A.K. Murchie. 2003. The impact of fungicides to control apple scab (*Venturia inaequalis*) on the predatory mite *Anystis baccarum* and its prey *Aculus schlechtendali* (apple rust mite) in Northern Ireland Bramley orchards. *Crop Protection* 22:1125–1130.
- Datnoff, L.E., and R.T. Nagata. 1999. Influence of Silicon on Gray Leaf Spot Development in St. Augustinegrass. *Phytopathology* 89:S19.
- Datnoff, L.E., F.A. Rodrigues, and K.W. Seebold. 2007. Silicon and plant disease. pp. 233–246 In: L.E. Datnoff, W.H. Elmer, and D.M. Huber (Eds.). *Mineral Nutrition and Plant Disease*. The American Phytopathological Society, St. Paul, Minnesota, U.S.
- Datnoff, L.E., G.H. Snyder, R.N. Raid, and D.B. Jones. 1991. Effect of calcium silicate on blast and brown spot intensities and yields of rice. *Plant Disease* 75:729–732.
- Elliott, C.L., and G.H. Snyder. 1991. Autoclave-induced digestion for the colorimetric determination of silicon in rice straw. *Journal of Agricultural Food Chemistry* 39:1118–1119.
- Facchini, P.J. 2001. Alkaloid biosynthesis in plants: Biochemistry, cell biology, molecular regulation, and metabolic engineering applications. *Annual Review of Plant Physiology and Plant Molecular Biology* 52:29–66.
- Fauteux, F., R.B. Wilfried, J.G. Menzies, and R.R. Belanger. 2005. Silicon and plant disease resistance against pathogenic fungi. *FEMS Microbiology Letters* 249:1–6.
- Fobert, P.R., and C. Després. 2005. Redox control of systemic acquired resistance. *Current Opinion Plant Biology* 8:378–382.
- Garbelotto, M., D.J. Schmidt, and T.Y. Harnik. 2007. Phosphite injections and bark application of phosphite + pentrabark™ control sudden oak death in coast. *Arboriculture & Urban Forestry* 33:309–316.
- Gozzo, F. 2003. Systemic acquired resistance in crop protection: From nature to a chemical approach. *Journal Agriculture Food Chemistry* 51:4487–4503.
- Hailey, L.E., and G.C. Percival. 2014. Comparative assessment of phosphites formulations for apple scab (*Venturia inaequalis*) control. *Arboriculture & Urban Forestry* 40:237–243.
- Hamel, S.C., and J. R. Heckman. 2000. Impact of mineral silicon products on powdery mildew in greenhouse grown turf. pp. 215–219. In: A.B. Gould (Ed.). *Rutgers Turfgrass Proceeding*. Center for Turfgrass Science, Rutgers University, New Brunswick, New Jersey, U.S.
- Hammerschmidt, R. 2003. Host cell death and the onset of systemic acquired resistance: Revisiting an old observation. *Physiology Molecular Plant Pathology* 62:315–316



- Hayasaka, T., H. Fujii, and K. Ishiguro. 2008. The role of silicon in preventing appressorial penetration by the rice blast fungus. *Phytopathology* 98:1038–1044.
- Ilhan, K., U. Arslan, and O.A. Karabulut. 2006. The effect of sodium bicarbonate alone or in combination with a reduced dose of tebuconazole on the control of apple scab. *Crop Protection* 25:963–967.
- Jamar, L., M. Cavelier, and M. Lateur. 2010. Primary scab control using a “during-infection” spray timing and the effect on fruit quality and yield in organic apple production. *Biotechnology Agronomy Society Environment* 14:423–439.
- Keeling, C.L., and J. Bohlmann. 2006. Genes, enzymes, and chemicals of terpenoid diversity in the constitutive and induced defense of conifers against insects and pathogens. *New Phytologist* 170:657–675.
- Kim, S.G., K.W. Kim, E.W. Park, and D. Choi. 2002. Silicon-induced cell wall fortification of rice leaves: A possible cellular mechanism of enhanced host resistance to blast. *Phytopathology* 92:1095–1103.
- Liang, Y.C., W.C. Sun, J. Si, and V. Römheld. 2005. Effect of foliar and root-applied silicon on the enhancement of induced resistance in *Cucumis sativus* to powdery mildew. *Plant Pathology* 54:678–685.
- Lichtenthaler, H.K., and A.R. Wellburn. 1983. Determinations of total carotenoids and chlorophylls a and b of leaf extracts in different solvents. *Biochemical Society Transactions* 11:591–593.
- Percival, G.C. 2016. The influence of systemic inducing agents on horse chestnut leaf miner (*Cameraria ohridella*) severity in white flowering horse chestnut (*Aesculus hippocastanum* L.). *Urban Forestry & Urban Greening* 20:97–102.
- Percival, G.C., and I. Haynes. 2008. The influence of systemic inducing resistance chemicals for the control of oak powdery mildew (*Microspora alphitoides*) applied as a therapeutic treatment. *Arboriculture & Urban Forestry* 34:191–200.
- Percival, G.C., K. Noviss, and I. Haynes. 2009. Field evaluation of systemic inducing resistance chemicals at different growth stages for the control of apple (*Venturia inaequalis*) and pear (*V. pirina*) scab. *Crop Protection* 28:629–633.
- Resende, R.S., F.Á. Rodrigues, R.V. Costa, and D.D. Silva. 2012. Silicon and fungicide effects on anthracnose in moderately resistant and susceptible sorghum lines. *Journal of Phytopathology* 161(1):11–17.
- Rodrigues, F.Á., N. Benhamou, L.E. Datnoff, J.B. Jones, and R.R. Bélanger. 2003. Ultrastructural and cytochemical aspects of silicon-mediated rice blast resistance. *Phytopathology* 93:535–546.
- Rondeau, E. 2001. Effect of potassium silicates on disease tolerance of bentgrass. *Seminaire de fin d'études, Centre de Recherche en Horticulture, Université Laval, Québec, Canada.*
- Saigusa, M., K. Onozawa, H. Watanabe, and K. Shibuya. 2000. Effects of porous hydrate calcium silicate on the wear resistance, insect resistance, and disease tolerance of turf grass “Miyako.” *Grassland Science*. 45:416–420.
- Schmidt, R.E., X. Zhang, and D.R. Chalmers. 1999. Response of photosynthesis and superoxide dismutase to silica applied to creeping bentgrass grown under two fertility levels. *Journal Plant Nutrition* 22:1763–1773.
- Schnabel, G., and L. Parisi. 1997. Sensitivity of *Venturia inaequalis* to five DMI fungicides, including the new triazole fluquinconazole, and to pyrimethanil. *Journal Plant Disease Protection* 104:36–46.
- Seebold, K.W., L.E. Datnoff, F.J. Correa-Victoria, T.A. Kucharek, and G.H. Snyder. 2004. Effects of silicon and fungicides on the control of leaf and neck blast in upland rice. *Plant Disease* 88:253–258.
- Sparla, F., L. Rotino, M.C. Valgimigli, P. Pupillo, and P. Trost. 2004. Systemic resistance induced by benzothiadiazole in pear inoculated with the agent of fire blight (*Erwinia amylovora*). *Scientia Horticulturae* 101:269–279.
- Sun, W.C., J. Zhang, Q.H. Fan, G.F. Xu, and Z.L. Li. 2010. Silicon-enhanced resistance to rice blast is attributed to silicon-mediated defense resistance and its role as physical barrier. *European Journal Plant Pathology* 128:39–49.
- Swait, A.A.J., and D.J. Butt. 1990. Fungicides as antisporengium against apple powdery mildew and scab. *Tests of Agrochemicals and Cultivars* 11. *Annals Applied Biology* (supplement) 116:36–37.
- Uriarte, R.F., H.D. Shew, and D.C. Bowman. 2004. Effect of soluble silica on brown patch and dollar spot of creeping bentgrass. *Journal Plant Nutrition* 27:325–339.
- Vallad, G.E., and R.M. Goodman. 2004. Systemic acquired resistance and induced systemic resistance in conventional agriculture. *Crop Science* 44:1920–1934.
- Villalta, O.N., W.S. Washington, and G. McGregor. 2004. Susceptibility of European and Asian pears to pear scab. *Plant Protection Quarterly* 19:2–4.
- Walters, D. 2009. Are plants in the field already induced? Implications for practical disease control. *Crop Protection* 28:459–465.
- Witzell, J., and J.A. Martin. 2008. Phenolic metabolites in the resistance of northern forest trees to pathogens – past experiences and future prospects. *Canadian Journal of Forest Research* 38:2711–2727.
- Zhang, G.L., D.Q. Gen, and H.C. Zhang. 2006. Silicon application enhances resistance to sheath blight (*Rhizoctonia solani*) in rice. *Journal Plant Physiology Molecular* 32:600–606.

Glynn C. Percival  
 R.A. Bartlett Tree Research Laboratory  
 Cutbush Lane  
 Shinfield, University of Reading  
 Reading, RG2 9AF UK  
 gpercival@bartlettuk.com

**Résumé.** Deux essais sur le terrain furent effectués avec des pommiers (*Malus* cv. Golden Delicious) et des poiriers (*Pyrus communis* 'Williams' Bon Chrétien') déjà établis afin d'évaluer l'efficacité d'un éventail de fertilisants à base de silicium (Si), d'un agent (dérivant de l'acide salicylique) induisant une résistance (IR) et d'une combinaison de fertilisants de silicium et d'IR lorsqu'appliqué à quatre différents stades de développement des arbres fruitiers (c'est-à-dire, au moment du débourrement, lorsque les feuilles sont encore en grappe, lorsque 90% des pétales sont tombés et au moment où les fruits apparaissent) contre des galles attaquant le feuillage des pommiers (*Venturia inaequalis*) et des poiriers (*V. pirina*). En parallèle, une évaluation comparative avec un fongicide synthétique conventionnel (penconazole) utilisé au Royaume-Uni pour le contrôle des galles sur les pommiers et les poiriers fut réalisée. La meilleure protection contre les galles de pommiers et de poiriers fut obtenue avec la combinaison d'un fertilisant à base de silicium et de l'agent inducteur IR ainsi qu'avec le fongicide synthétique penconazole alors que les résultats des essais en 2014 et en 2015 furent statistiquement comparables sur le niveau de contrôle obtenu. L'efficacité des composés prévenant les galles (c'est-à-dire la réduction des dommages causés par les galles aux feuilles et aux fruits), la quantité plus élevée de chlorophylle dans les feuilles et la récolte fruitière accrue fut confirmée lorsque des fertilisants de silicium furent appliqués à chacun des quatre stades de développement durant deux saisons de croissance; cependant, il y avait peu de différences dans le niveau de contrôle observé selon les différents fertilisants de silicium. L'agent inducteur IR utilisé seul s'est avéré le moins efficace des composés.

**Zusammenfassung.** Es wurden zwei Feldversuche an etablierten Äpfeln (*Malus* cv. Golden Delicious) und Birnen (*Pyrus communis* 'Williams' Bon Chrétien') durchgeführt, um die Effizienz gegen den Befall mit Blattkrankheiten an Apfel (*Venturia inaequalis*) und Birne (*V. pirina*) mit einer Auswahl von Silizium (Si)-Düngern, einem auf Salicylsäure basierendem Agens zur Einführung von Resistenz (IR) und einer Kombination aus Si+IR, die in vier verschiedenen Wachstumsstadien der Baumentwicklung (z. B. Knospenöffnung, Blattaustrieb, 90% der Blüten gefallen, frühe kleine Früchte) aufgetragen wurden, zu untersuchen. Zusätzlich wurde eine vergleichen-

de Evaluation mit einem konventionellen synthetischen Fungizid (Penconazol), welches in Großbritannien gegen Apfel- und Birnenschorf eingesetzt wird, durchgeführt. Der größte Schutz gegen Apfel- und Birnenschorf wurde durch eine Kombination mit einem Si-Dünger+IR-Agens und dem synthetischen Fungizid Penconazol in den beiden Feldversuchen in 2014 und 2015 erzielt, wo statistisch vergleichbare Grade an Kontrolle erzielt wurden. Die Effizienz der vor Schorfbefall schützenden Verbindung (z.B. reduzierte Schorfbefall an Blättern und Früchten) erhöhte den Blattchlorophyllanteil und Fruchtertrag und wurde bestätigt, wenn jeder der Si-Dünger während der zwei Vegetationsperioden in den vier Wachstumsphasen appliziert wurde. Dennoch gab es kleine Differenzen beim Ausmaß der Kontrolle, die von jedem Si-Dünger gewährt wurde. Das allein verwendete IR-Agens zeigte die geringste Effizienz bei der Behandlung gegen Schorf.

**Resumen.** Se realizaron dos ensayos de campo usando plantas establecidas de manzana (*Malus* cv. Golden Delicious) y pera (*Pyrus communis* 'Williams' Bon Chrétien ') para evaluar la eficacia de una gama de fertilizantes de silicio (Si), un agente inductor de resistencia (IR) basado en ácido salicílico y una combinación de Si + IR cuando es aplicado en cuatro etapas de crecimiento del árbol (brote, racimo verde, 90% caída de pétalos, fruto temprano) contra los patógenos foliares de la manzana (*Venturia inaequalis*) y antracnosis de la pera (*V. pirina*). Además, se realizó una evaluación comparativa de un fungicida sintético convencional (penconazol) utilizado en el Reino Unido para el control de la antracnosis de la manzana y pera. La mayor protección contra la antracnosis de la manzana y la pera provino de una combinación de un fertilizante de Si + agente de IR y el fungicida sintético penconazol en los ensayos de campo 2014 y 2015 donde se lograron grados de control estadísticamente comparables. La eficacia como compuestos protectores de antracnosis (es decir, reducción de la severidad en hojas y frutos) mejoró el contenido de clorofila de la hoja y el rendimiento de fruta, y se confirmó cuando cada fertilizante de Si se aplicó en cuatro etapas de crecimiento durante dos temporadas; sin embargo, hubo poca diferencia en el grado de control conferido por cada fertilizante de Si. El agente de IR usado solo demostró ser el compuesto protector de antracnosis menos efectivo.