breakage, elimination of branch stubs, pruning of shaded or weakened branches, removal of plant debris, and disinfection of tools and equipment will help reduce the likelihood of spread and infection.

## **Literature Cited**

- Baker, K.F. and R.J. Cook. 1973. Biological Control of Plant Pathogens. W.H. Freeman, San Francisco.
- Burke, M.J., L.V. Gusta, H.A. Quamme, C.J. Weiser, and P.H. Li. 1976. Freezing and injury in plants. Annu. Rev. Plant Physiol. 27:507-528.
- Colhoun, J. 1973. Effects of environmental factors on plant disease. Annu. Rev. Phytopathol. 11:343-364.
- Crist, C.R. and D.F. Schoeneweiss. 1975. The influence of controlled stresses on susceptibility of European white birch stems to attack by Botryosphaeria dothidea. Phytopathology 65:369-373.
- Levitt, J. 1972. Responses of Plants to Environmental Stresses. Academic Press, New York.
- Li, P.H. 1978. Plant cold hardiness research. HortScience 13:222-224.
- Schoeneweiss, D.F. 1974. Tubercularia ulmea canker of tallhedge: influence of freezing stress on disease susceptibility. Plant Dis. Reptr. 58:937-941.

- 8. Schoeneweiss, D.F. 1975. Predisposition, stress, and plant disease. Annu. Rev. Phytopathol. 13:193-211.
- Schoeneweiss, D.F. 1975. A method for controlling plant water potentials for studies on the influence of water stress on disease susceptibility. Can. J. Bot. 53:647-652.
- Schoeneweiss, D.F. 1978. Water stress as a predisposing factor in plant disease. p. 61-99. In "Water Deficits and Plant Growth," Vol. V: T.T. Kozlowski, ed. Academic Press, New York.
- Schoeneweiss, D.F. 1978. The influence of stress on diseases of nursery and landscape plants. J. Arboric. 4:217-225.
- 12. Schoeneweiss, D.F. and E.G. Wene. 1977. Freezing stress predisposes Euonymus alatus to attack by Nectria cinnabarina. Plant Dis. Reptr. 61:921-925.
- Wene, E.G. 1979. Stress predisposition of woody plants to Botryosphaeria dothidea stem canker. Ph.D. Thesis, University of Illinois.

Plant Pathologist Section of Botany and Plant Pathology Illinois Natural History Survey Urbana, Illinois

**ABSTRACTS** 

Haller, J.M. 1979. Galls. Am. Forests 85(12): 12-15, 51-53.

Galls are globular, semiglobular,, cylindrical, or disklike swellings produced on leaf blades, leaf petioles, or twigs by the sting of parasitic wasps; by midges, aphids, or other insects; or by the infestation of bacteria or fungi. Although each gall is characteristic of the insect or bacterium or fungus that causes it, the gall itself is actually part of the plant. It is composed of plant tissues diverted from their normal development and forced to serve the ends of the parasite. The gall is alive only so long as the plant is alive. The great majority of shade-tree galls are caused by egg-laying insects, especially tiny wasps. Uninformed homeowners discovering galls for the first time often take them as symptomatic of a dangerous disease and may resort to drastic remedies. Felling and topping are, of course, useless, since the gall-producing insects may fly to any tree from any point of the compass. Spraying is similarly of very doubtful value.

Shurtleff, M.C. 1980. Fungicides for ornamentals — selection and use. Grounds Maintenance 15(1): 20-21, 24, 82-83, 86.

Fungicides are available to protect plant seeds, foliage, flowers, fruits and roots from infection, but no single fungicide is suitable against all types of disease-causing fungi. Because most fungicides in general use have protective-contact action, they must uniformly coat all susceptible plant parts before fungus invasion occurs. Systemic fungicides, however, are absorbed and translocated within the plant to control a disease. In general, fungicide sprays, dusts, dips and soil drenches will not check or eradicate disease that is well established or protect against diseases caused by bacteria, mycoplasmas, spiroplasmas, viruses, viroids, nematodes or parasitic plants. See the chart and tables for information on fungicide efficacy for various categories of diseases and lists of generic (common) names followed by their trade names.